

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets

(11)

EP 0 903 343 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:

24.03.1999 Bulletin 1999/12

(21) Application number: 98117153.1

(22) Date of filing: 10.09.1998

(51) Int. Cl.⁶: C07D 237/32, C07D 239/88,
C07D 265/22, C07D 401/04,
C07D 239/96, C07D 243/08,
C07D 473/10, A61K 31/495

(84) Designated Contracting States:

AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE

Designated Extension States:

AL LT LV MK RO SI

(30) Priority: 19.09.1997 JP 255025/97

(71) Applicant: SSP Co., Ltd.
Chuo-ku, Tokyo (JP)

(72) Inventors:


- Nagao, Yoshihiro
Narita-shi, Chiba (JP)
- Konno, Fujiko
Inga-bun, Chiba (JP)

- Kotake, Jiro
Ichikawa-shi, Chiba (JP)
- Ishii, Fumio
Sendai-shi, Miyagi (JP)
- Honda, Haruyoshi
Inba-gun, Chiba (JP)
- Sato, Susumu
Narita-shi, Chiba (JP)

(74) Representative:

Hartz, Nikolai F., Dr. et al
Wächtershäuser & Hartz,
Patentanwälte,
Tal 29
80331 München (DE)

(54) Alpha-Substituted phenylpropionic acid derivative and medicine containing the same

(57) Disclosed herein are an α -substituted phenylpropionic acid derivative represented by the following general formula (1):

(1)

09/152,785

wherein W is a (substituted) lactam ring, A is an alkylene or alkyleneoxy group or the like, X is O, S, NH or CH₂, Y¹ is an amino, hydroxy or lower alkoxy group, R¹ is H, alkyl group or the like, R² is an alkyl or phenyl group or the like, and R³ is H, alkyl or alkoxy group or the like, or a salt thereof, and a medicine comprising such a compound as an active ingredient. The compound is excellent in the effect of lowering blood glucose and lipid.

EP 0 903 343 A1

BEST AVAILABLE COPY

Description

BACKGROUND OF THE INVENTION

5 Field of the Invention:

[0001] The present invention relates to α -substituted phenylpropionic acid derivatives or salts thereof, which are excellent in the effect of lowering blood glucose and lipid, and medicines comprising such a compound as an active ingredient.

10

Description of the Background Art:

[0002] Non-insulin dependent diabetes mellitus (NIDDM) is a disease caused by insulin resistance in a target tissue of insulin and impaired insulin secretion from β cells of pancreas. Sulfonylureas and insulin, which are in wide use for treating NIDDM at present, are agents for mainly improving impaired insulin secretion. Of these, the sulfonylureas, which are oral drugs, have a strong antidiabetic effect based on both pancreatic action and extrapancreatic action, but often cause serious hypoglycemia. Therefore, care must be taken in using them.

15

[0003] In recent years, the importance of insulin resistance in NIDDM has come to be indicated, so that there has been a demand for development of a drug exhibiting an antidiabetic effect by lightening insulin resistance in a target tissue of insulin without stimulating insulin secretion. As compounds having such an effect, have been developed thiazolidine derivatives such as troglitazone and pioglitazone (Japanese Patent Application Laid-Open Nos. 22636/1980, 51189/1985 and 157522/1994, etc.). In addition, some thiazolidine derivatives having a like effect and a bicyclic lactam structure or cyclic urethane structure have also been reported [Japanese Patent Application (KOHYO) Nos. 502144/1994, 502145/1994 and 502146/1994 (through PCT route), etc.]. Further, a great number of derivatives having no thiazolidine ring have also been reported [for example, Japanese Patent Application Laid-Open No. 170478/1991, Japanese Patent Application (KOHYO) No. 508054/1993 (through PCT route), etc.]. Furthermore, arylpropionic acid derivatives have also been reported as antidiabetics (WO91/19702, Japanese Patent Application Laid-Open Nos. 325250/1996, 325263/1996 and 325264/1996, etc.).

20

25

[0004] However, the blood glucose-lowering effect brought about by these compounds which lighten insulin resistance has been not yet sufficient.

30

[0005] On the other hand, hyperlipemia and obesity have been becoming a problem in modern times when satiation and lack of exercise have been chronic, and there is hence a demand for development of medicines for treating these diseases.

35 SUMMARY OF THE INVENTION


[0006] Accordingly, it is an object of the present invention to provide a novel compound excellent in the effect of lowering blood glucose and lipid.

[0007] With the foregoing circumstances in view, the present inventors have synthesized various kinds of compounds and carried out an extensive investigation as to their pharmacological effects. As a result, it has been found that α -substituted phenylpropionic acid derivatives represented by a general formula (1), which will be described subsequently, are excellent in the effect of lowering blood glucose and lipid and hence are useful in preventing or treating diabetes mellitus, hyperlipemia, obesity and the like, thus leading to completion of the present invention.

40

[0008] According to the present invention, there is thus provided an α -substituted phenylpropionic acid derivative represented by the following general formula (1):

45



50

55 wherein W is a monocyclic or bicyclic lactam ring which may be substituted, A is an alkylene, alkyleneoxy or alkylene-carbonyl group which may be substituted by at least one hydroxy group, X is O, S, NH or CH₂, Y¹ is an amino, hydroxyamino, hydroxyalkylamino, monoalkylamino, dialkylamino, cyclic amino, hydroxy or lower alkoxy group, R¹ is a hydrogen atom, lower alkyl group, hydroxyalkyl group, alkoxyalkyl group, halogenoalkyl group or COY² (in which Y² is

55

an amino, hydroxyamino, hydroxyalkylamino, monoalkylamino, dialkylamino, cyclic amino, hydroxy or lower alkoxy group), R^2 is a lower alkyl, hydroxyalkyl, alkoxyalkyl or halogenoalkyl group, COY^2 (in which Y^2 has the same meaning as defined above), or a phenyl, pyridyl or aralkyl group which may be substituted, and R^3 is a hydrogen or halogen atom, or an alkyl, alkoxy, halogenoalkyl, amino, hydroxy or acyl group, or a salt thereof.

[0009] According to the present invention, there is also provided a medicine comprising the α -substituted phenylpropionic acid derivative represented by the general formula (1) or the salt thereof as an active ingredient.

[0010] According to the present invention, there is further provided a medicinal composition comprising the α -substituted phenylpropionic acid derivative represented by the general formula (1) or the salt thereof and a pharmaceutically acceptable carrier.

[0011] According to the present invention, there is still further provided use of the α -substituted phenylpropionic acid derivative represented by the general formula (1) or the salt thereof for a medicine.


[0012] According to the present invention, there is yet still further provided a method of treating diabetes mellitus and/or hyperlipemia, which comprises administering the α -substituted phenylpropionic acid derivative represented by the general formula (1) or the salt thereof to a patient.

[0013] The compounds (1) according to the present invention are excellent in the effect of lowering blood glucose and lipid and are hence useful as agents for preventing or treating diabetes mellitus, hyperlipemia, obesity and the like.

[0014] The above and other objects, features and advantages of the present invention will be readily appreciated as the same becomes better understood from the preferred embodiments of the present invention, which will be described subsequently in detail, and from the appended claims.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

[0015] The lactam ring represented by W in the α -substituted phenylpropionic acid derivatives according to the present invention, which are represented by the general formula (1), is preferably selected from among, for example, groups represented by the following (W-1) to (W-9):



wherein R^4 is a hydrogen or halogen atom, an alkyl, alkoxy, halogenoalkyl, amino, hydroxy, cyano, carbamoyl, acyl, nitro, carboxy or sulfonamide group, or a phenyl or benzyloxy which may be substituted, R^5 is a hydrogen atom, an alkyl

group, or an aryl, aralkyl or pyridyl group which may be substituted, R^6 is a hydrogen atom or a lower alkyl group, R^7 is a lower alkyl, phenyl or aralkyl group, Z^1 is O, S, CH_2 or NR^5 (in which R^5 has the same meaning as defined above), Z^2 is N or CH, and m is an integer of 1 to 4.

[0016] In the above formulae, the alkyl groups represented by R^4 and R^5 are preferably linear or branched alkyl groups having 1 to 8 carbon atoms. Examples thereof include methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, tert-butyl, pentyl, hexyl, heptyl and octyl groups. The alkoxy group represented by R^4 is preferably a linear or branched alkoxy group having 1 to 8 carbon atoms. Examples thereof include methoxy, ethoxy, propoxy, butoxy, pentyloxy, hexyloxy, heptyloxy and octyloxy groups. Examples of the halogen atom include fluorine, chlorine, bromine and iodine atoms. The halogenoalkyl group is preferably an alkyl group substituted by 1 to 3 halogen atoms and having 1 to 8 carbon atoms. Examples thereof include trifluoromethyl, trichloromethyl and tribromomethyl groups. Examples of the acyl group include alkanoyl groups having 1 to 9 carbon atoms, such as formyl, acetyl and propionyl groups, and aroyl groups such as a benzoyl group. Examples of substituents on the phenyl or benzyloxy group include the alkyl groups, alkoxy groups, halogen atoms, halogenoalkyl groups and acyl group, which have been mentioned above, and besides amino, hydroxy, cyano, carbamoyl, nitro, carboxy, sulfonamide, phenyl and benzyloxy groups.

[0017] The lower alkyl groups represented by R^6 and R^7 are preferably linear or branched alkyl groups having 1 to 6 carbon atoms. Examples thereof include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl and sec-butyl groups. Examples of the aryl group represented by R^5 include aryl groups having 6 to 14 carbon atoms, such as phenyl and naphthyl groups. Examples of the aralkyl groups represented by R^5 and R^7 include phenyl- C_{1-6} -alkyl groups such as benzyl and phenethyl groups. Examples of substituents on the aryl, aralkyl or pyridyl group represented by R^5 include the same substituents as those mentioned in the phenyl group represented by R^4 . The group W is particularly preferably (W-1), (W-2) (W-4) or (W-5).

[0018] The group A in the general formula (1) is preferably a linear or branched alkylene group having 1 to 8 carbon atoms, which may be substituted by 1 to 5 hydroxy groups, a linear or branched alkyleneoxy group having 1 to 8 carbon atoms, which may be substituted by 1 to 5 hydroxy groups, or a linear or branched alkylene carbonyl group having 2 to 9 carbon atoms, which may be substituted by 1 to 5 hydroxy groups. Specific examples of A include ethylene, trimethylene, propylene, tetramethylene, butylene, ethyleneoxy, trimethyleneoxy, 2-hydroxytrimethyleneoxy, propyleneoxy, butyleneoxy, methylenecarbonyl, ethylenecarbonyl and trimethylenecarbonyl groups. Of these, the ethylene, trimethylene, ethyleneoxy and 2-hydroxytrimethyleneoxy groups are particularly preferred.

[0019] An atomic group represented by X is preferably an oxygen atom.

[0020] With respect to the groups represented by Y^1 and Y^2 , the alkyl moieties of the mono- or dialkylamino groups or hydroxyalkylamino groups are linear or branched alkyl groups having 1 to 6 carbon atoms. Examples thereof include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl and tert-butyl groups. Examples of the cyclic amino groups include 4-membered to 7-membered cyclic amino groups such as piperazinyl, piperidinyl, pyrrolidinyl and azetidiny groups. Examples of the alkoxy groups include linear or branched alkoxy groups having 1 to 6 carbon atoms, such as methoxy, ethoxy, propoxy and butoxy groups.

[0021] Examples of the lower alkyl groups represented by R^1 and R^2 include linear or branched alkyl groups having 1 to 6 carbon atoms, such as methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl and tert-butyl groups.

[0022] The hydroxyalkyl group represented by R^2 is preferably a hydroxy- C_{1-6} -alkyl group. Examples thereof include hydroxyethyl and hydroxypropyl groups. The alkoxyalkyl group represented by R^2 is preferably a C_{1-6} -alkoxy- C_{1-6} -alkyl group. Examples thereof include methoxyethyl, ethoxyethyl and methoxypropyl groups. The halogenoalkyl group represented by R^2 is preferably an alkyl group substituted by 1 to 3 halogen atoms and having 1 to 6 carbon atoms. Examples thereof include trifluoromethyl and trifluoroethyl groups.

[0023] Examples of the aralkyl group represented by R^2 include phenyl- C_{1-6} -alkyl groups such as benzyl and phenethyl groups. Examples of substituents on the phenyl, pyridyl or aralkyl group represented by R^2 include alkyl groups, alkoxy groups, halogen atoms, halogenoalkyl groups, acyl groups, amino group, hydroxy group, cyano group, carbamoyl group, nitro group, carboxy group, sulfonamide group, phenyl group and benzyloxy group. R^1 is preferably a hydrogen atom. R^2 is preferably a lower alkyl group.

[0024] The alkyl group represented by R^3 is preferably a linear or branched alkyl group having 1 to 8 carbon atoms. Examples thereof include methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, tert-butyl, pentyl, hexyl, heptyl and octyl groups. The alkoxy group represented by R^3 is preferably a linear or branched alkoxy group having 1 to 8 carbon atoms. Examples thereof include methoxy, ethoxy, propoxy, butoxy, pentyloxy, hexyloxy, heptyloxy and octyloxy groups. Examples of the halogen atom include fluorine, chlorine, bromine and iodine atoms. The halogenoalkyl group is preferably an alkyl group substituted by 1 to 3 halogen atoms and having 1 to 8 carbon atoms. Examples thereof include trifluoromethyl, trichloromethyl and tribromomethyl groups. Examples of the acyl group include alkanoyl groups having 1 to 9 carbon atoms, such as formyl, acetyl and propionyl groups, and aroyl groups such as a benzoyl group. R^3 is preferably a hydrogen atom.

[0025] The compounds (1) according to the present invention may include various kinds of pharmaceutically acceptable solvates such as hydrates, and those of polymorphism. Since the compounds (1) according to the present inven-


tion have an asymmetric carbon atom, stereoisomers exist. All these isomers are included in the present invention.

[0026] No particular limitation is imposed on salts of the compounds (1) according to the present invention so far as they are pharmaceutically acceptable salts. Preferable examples of such salts include hydrogen halide salts such as hydrofluorides, hydrochlorides, hydrobromides and hydriodides; inorganic acid salts such as carbonates, nitrates, perchlorates, sulfates and phosphates; lower alkyl-sulfonates such as methanesulfonates, ethanesulfonates and trifluoromethanesulfonates; arylsulfonates such as benzenesulfonates and p-toluenesulfonates; organic acid salts such as fumarates, maleates, succinates, citrates, tartrates and oxalates; amino acid salts such as glutamates and aspartates; and salts with alkali and alkaline earth metals such as sodium, potassium and calcium.

[0027] The compounds (1) according to the present invention can be prepared, for example, in accordance with any of the following Preparation Processes 1 to 8.

<Preparation Process 1>

Case of $R^1 = H$ and $X = O$ in the general formula (1):



wherein R^2 , R^3 , W and A have the same meanings as defined above, A^2 is an alkyleneoxy group which may be substi-

tuted by at least one hydroxyl group, Y^3 is an amino, hydroxyl or lower alkyl group, Q^1 is a leaving group, and Q^2 is a halogen atom.

[0028] More specifically, a compound (1-1) according to the present invention is prepared in the following manner. A compound represented by a general formula (2) is reacted with a compound represented by a general formula W-H to form a compound represented by a general formula (5) (Step 1). The 4-substitution product of compounds (5) may also be synthesized by the reaction of a compound represented by a general formula (3) with a compound represented by a general formula (4) (Step 2). The compound (5) is then subjected to a Wittig reaction with (methoxymethyl)triphenylphosphonium chloride (Step 3), and the resultant compound represented by a general formula (6) is reacted with its corresponding alcohol to form an acetal derivative represented by a general formula (7) (Step 4). The acetal derivative is then reacted with trimethylsilyl nitrile to form a compound represented by a general formula (8) (Step 5). Finally, the compound (8) is hydrolyzed or reacted with an alcohol in the presence of an acid catalyst (Step 6), whereby the compound (1-1) according to the present invention can be prepared. The individual steps will hereinafter be described in detail.

Step 1:

[0029] The compound (2) is reacted with the compound W-H in the presence of proper base and solvent, whereby the compound (5) can be obtained.

[0030] The compound (2) can be prepared by halogenation or sulfonylation of the terminal hydroxy group in the benzaldehyde derivative substituted by a hydroxyalkoxy group or hydroxyalkyl group, which has been purchased as a commercially available reagent or synthesized in accordance with a known method [for example, a method described in Journal of Heterocyclic Chemistry, 6, 243 (1969) or Japanese Patent Application Laid-Open No. 92228/1996]. Examples of the leaving group (Q^1) thereof include halogen atoms such as chlorine, bromine and iodine atoms, and methanesulfonyloxy, p-toluenesulfonyloxy and trifluoromethanesulfonyloxy groups. Of these, the methanesulfonyloxy group is particularly preferred.

[0031] Examples of the base used in the reaction of the compound (2) with the compound W-H include sodium hydride, calcium hydride, potassium t-butoxide, sodium hydroxide, potassium hydroxide and potassium carbonate. No particular limitation is imposed on the solvent used herein so far as it does not affect the reaction. Examples of the solvent used include ethers such as tetrahydrofuran and dioxane; hydrocarbons such as benzene and toluene; amides such as dimethylformamide, dimethylacetamide and N-methyl- α -pyrrolidone; and sulfoxides such as dimethyl sulfoxide. The reaction is carried out in a temperature range of from a temperature under ice cooling to a reflux temperature under heating. It is particularly preferred to conduct the reaction by heating and stirring the reactants at 70 to 100°C for 2 to 5 hours in the presence of potassium carbonate in dimethylformamide.

Step 2:

[0032] The compound (3) which is a starting material can be prepared, for example, in accordance with the method described in Journal of Medicinal Chemistry, 11, 1038 (1968); or Journal of Medicinal Chemistry, 38, 130 (1995). This compound (3) is reacted with the p-halogenobenzaldehyde derivative (4) in the presence of proper base and solvent, whereby the compound (5) can be obtained. Examples of the halogen atom Q^2 in the p-halogenobenzaldehyde derivative (4) include, fluorine, chlorine, bromine and iodine atoms, with the fluorine atom being particularly preferred. Examples of the base used in this reaction include sodium hydride, potassium t-butoxide, sodium hydroxide and potassium hydroxide. No particular limitation is imposed on the solvent used herein so far as it does not affect the reaction. Examples of the solvent used include ethers such as tetrahydrofuran and dioxane; hydrocarbons such as benzene and toluene; amides such as dimethylformamide, dimethylacetamide and N-methyl- α -pyrrolidone; and sulfoxides such as dimethyl sulfoxide. The reaction is carried out in a temperature range of from a temperature under ice cooling to a reflux temperature under heating. The reaction time is about 0.5 to 24 hours. It is particularly preferred to conduct the reaction by adding sodium hydride little by little to a dimethyl sulfoxide solution of a mixture of the compound (3) and the compound (4) under ice cooling and then stirring the resultant mixture for 1 to 3 hours at room temperature or so.

Step 3:

[0033] The compound (5) is subjected to the Wittig reaction with (methoxymethyl)triphenylphosphonium chloride, which is a commercially available reagent, in the presence of proper base and solvent, whereby the compound (6) can be obtained [the compound (6) is obtained as a mixture of E:Z]. Examples of the base used in this reaction include n-butyllithium, sec-butyllithium, lithium diisopropylamide, potassium t-butoxide and sodium methoxide. No particular limitation is imposed on the solvent used herein so far as it does not affect the reaction. Examples thereof include ethers such as diethyl ether and tetrahydrofuran; hydrocarbons such as benzene and toluene; and alcohols such as ethanol.

It is particularly preferred to conduct the reaction by preparing lithium diisopropylamide in a flask containing tetrahydrofuran and stirring the reaction mixture at a reaction temperature of from -10°C to 20°C for about 3 to 5 hours.

Step 4:

[0034] The compound (6) is reacted with its corresponding alcohol in the presence of an acid catalyst, whereby the compound (7) can be prepared.

[0035] When the alcohol has a low boiling point, the alcohol itself is used as a solvent. When the alcohol has a high boiling point on the other hand, a hydrocarbon such as benzene or toluene, or an amide such as dimethylformamide or dimethylacetamide may be used as a solvent. Examples of the acid catalyst include p-toluenesulfonic acid and methanesulfonic acid. The reaction is carried out in a temperature range of from a warming temperature to a reflux temperature under heating. The reaction time is about 1 to 24 hours.


Step 5:

[0036] The compound (7) is reacted with an excess amount of trimethylsilyl nitrile in the presence of a boron trifluoride etherate catalyst in methylene chloride, whereby the compound (8) can be prepared. The reaction is carried out at a temperature of from 0 to 20°C for about 0.5 to 2 hours.

Step 6:

[0037] In general, the compound (8) is hydrolyzed under basic conditions, whereby the compound (1-1) according to the present invention can be prepared. Examples of the base used include sodium hydroxide and potassium hydroxide. As a solvent for the reaction, is used a mixed solvent of ethanol-water, dioxane-water or the like. The reaction is carried out in a temperature range of from 80°C to a reflux temperature for about 0.5 to 5 hours. An ester derivative corresponding to the compound (1-1) can be prepared by using a compound in which Y³ in the compound (1-1) according to the present invention is NH₂, and heating this compound and an alcohol with stirring in the presence of a catalytic amount of a Lewis acid such as titanium tetrachloride in 1N hydrochloric acid.

<Preparation Process 2> Case of $R^1 = H$ and $X = O$ in the general formula (1):



wherein W , R^2 , R^3 , Y^3 , A , A^2 and Q^1 have the same meanings as defined above, and A^1 means A or a single bond.

[0038] More specifically, a benzaldehyde derivative represented by a general formula (9) is allowed to react in the same manner as in Step 3 to Step 6 described in Preparation Process 1 to form a compound represented by a general formula (10). The benzyl group which is a protecting group is then removed from the compound (10) to obtain a com-

compound represented by a general formula (11) (Step 7). After the terminal hydroxy group of this compound is converted into a leaving group (Q^1) (Step 8), the resultant compound is allowed to react in the same manner as in Step 1 of Preparation Process 1, whereby the compound (1-1) according to the present invention can be prepared. When A^1 in the compound represented by the general formula (11) is a single bond, the compound is subjected to a condensation reaction with a compound represented by the general formula (3), whereby a compound (1-2) according to the present invention can be obtained (step 9). The individual steps will hereinafter be described.

Step 7:

[0039] The compound (9), which is a starting material, is purchased as a commercially available reagent or synthesized in accordance with a known method [for example, a method described in Journal of Heterocyclic Chemistry, 6, 243 (1969) or Japanese Patent Application Laid-Open No. 92228/1996], and is converted into a compound of the general formula (10) in the same manner as in Step 3 to Step 6 described in Preparation Process 1. This compound (10) is reduced by catalytical hydrogenation, whereby the compound of the general formula (11) can be prepared. Examples of the catalyst used in this reaction include palladium catalysts such as palladium on charcoal, palladium black and palladium hydroxide; platinum catalysts such as platinum oxide and platinum black; and nickel catalysts such as Raney nickel. No particular limitation is imposed on a solvent used so far as it does not affect the reaction. Examples thereof include methanol, ethanol, dioxane, dimethylformamide, acetic acid and a mixed solvent of ethanol-acetic acid. The reaction is carried out at atmospheric pressure or under pressure at room temperature or under heating at about 60 to 100°C.

Step 8:


[0040] The compound (12) is prepared by halogenating or sulfonylating the terminal hydroxy group of the phenylpropionic acid derivative (11), the benzene ring of which has been substituted by a hydroxyalkoxy group, hydroxyalkyl group or hydroxyalkanoyl group, in the presence or absence of a base and a solvent. Examples of the leaving group (Q^1) thereof include halogen atoms such as chlorine, bromine and iodine atoms, and methanesulfonyloxy, p-toluenesulfonyloxy and trifluoromethanesulfonyloxy groups. Of these, the methanesulfonyloxy group is particularly preferred. No particular limitation is imposed on the solvent used in the reaction so far as it does not affect the reaction. Examples of the solvent used include chlorinated hydrocarbons such as methylene chloride and chloroform; ethers such as tetrahydrofuran and dioxane; hydrocarbons such as benzene and toluene; amides such as dimethylformamide, dimethylacetamide and N-methyl- α -pyrrolidone; and sulfoxides such as dimethyl sulfoxide. Examples of the base used include triethylamine, pyridine, 4-dimethylaminopyridine, sodium hydride, sodium hydroxide and sodium carbonate.

[0041] The compound (12) thus obtained can be converted into the compound (1-1) according to the present invention in the same manner as in Step 1 of Preparation Process 1.

Step 9:

[0042] Among the compounds represented by the general formula (11), a compound in which A^1 is a single bond can be converted into the compound (1-2) according to the present invention by a Mitsunobu reaction (see Organic Reaction 42, 335) with a compound represented by the general formula (3), which has been described in Preparation Process 1. More specifically, each 1 to 3 equivalents of triphenylphosphine and dialkyl (dimethyl, diethyl or diisopropyl) azobiscarboxylate are added to a solution of the compound (3) and compound (11) in methylene chloride, tetrahydrofuran, benzene, toluene, ether, dioxane or dimethylformamide, and the reaction is conducted in a temperature range of from -5°C to a reflux temperature under heating for about 1 to 24 hours, whereby the compound (1-2) according to the present invention can be obtained.

<Preparation Process 3> Case of $R^1 = H$ in the general formula (1):



wherein W, A, R^2 , R^3 , A^1 , A^2 and Y^3 have the same meanings as defined above, and X^1 is O, S or CH_2 .

[0043] A compound represented by the general formula (9) is condensed with a compound represented by a general formula (13) to form a compound represented by a general formula (14) (Step 10). The compound thus obtained is allowed to react in the same manner as in Step 6 and Step 7 of Preparation Process 2, whereby a compound repre-

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000


5
10
15
20
25
30
35
40
45
50
55

sent by a general formula (16) can be obtained. Subsequently, the compound (16) can be converted into a compound (1-3) according to the present invention in the same manner as in Step 8 and Step 1 of Preparation Process 2. When A¹ in the compound represented by the general formula (16) is a single bond, the compound is subjected to a Mitsunobu reaction in the same manner as in Step 9 of Preparation Process 2, whereby a compound (1-4) according to the present invention can be obtained. Step 10 will hereinafter be described in detail.

Step 10:

[0044] The compound (13), which is a starting material, can be purchased as a commercially available reagent or prepared in accordance with a known method [for example, a method described in Organic Synthesis Collect, Volume II, 387, Japanese Patent Application Laid-Open No. 136391/1994, or the like]. This compound is condensed with the compound represented by the general formula (9) in the presence of a proper base or catalyst, whereby the compound (a mixture of E:Z) of the general formula (14) can be prepared. Examples of the base used in the reaction include sodium hydride, potassium t-butoxide and pyridine. When the catalyst is used, piperidine and acetic acid, piperidinium acetate, piperidinium benzoate, or the like may be used. No particular limitation is imposed on a solvent used so far as it does not affect the reaction. Examples of the solvent used include ethers such as tetrahydrofuran and dioxane; hydrocarbons such as benzene and toluene; amides such as dimethylformamide, dimethylacetamide and N-methyl- α -pyrrolidone; and sulfoxides such as dimethyl sulfoxide.

<Preparation Process 4> Case of $R^1 = H$ in the general formula (1):



wherein W , A , Y^1 , R^2 , R^3 , X^1 and A^2 have the same meanings as defined above, R^8 is a lower alkyl group, and X^2 is O or NH .

[0045] More specifically, a compound represented by a general formula (17) is converted into a compound (1-5)

Step 13:

[0049] The compound represented by the general formula (20) (EP 0787727-A1), which is a starting material, is hydrolyzed under basic conditions and then reacted with its corresponding halide, whereby the compound (1-7) according to the present invention can be prepared. Specifically, after the compound of the general formula (20) is gently heated to reflux for 0.5 to 1 hour in a 15% aqueous solution of sodium hydroxide, a methanol solution of the corresponding halide (iodide, bromide or chloride) is added at room temperature, and the mixture is stirred for about 1 to 3 hours, whereby the compound (1-7) according to the present invention can be obtained.

Step 14:

[0050] The carboxylic acid, which is the compound (1-7) according to the present invention, is converted into an active derivative (for example, an acid halide or mixed acid anhydride) and then reacted with any of various amines and alcohols, or the compound (1-7) according to the present invention is reacted with any of various amines and alcohols in the presence of a proper condensation agent, whereby the compound (1-8) according to the present invention can be prepared.

[0051] Examples of the condensation agent used include carbonyldiimidazole, 1-hydroxy-2(1H)-pyridone, N-hydroxysuccinimide, diphenylphosphorylazide, N,N-dicyclohexylcarbodiimide and 1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide hydrochloride. The reaction is carried out in the presence of a proper base, for example, an organic base such as triethylamine or pyridine, according to the kind of the condensation agent used.

according to the present invention by an ordinary alkylating method (Step 11). Subsequently, this compound is hydrolyzed or reacted with any of various amines, whereby a compound (1-6) according to the present invention can be prepared (Step 12). Alternatively, the compound (5) and a compound (18) are successively treated in the same manner as in Step 10 of Preparation Process 3 and Step 7 of Preparation Process 2, whereby the compound (1-5) according to the present invention can also be obtained. The individual steps will hereinafter be described.

Step 11:


[0046] The compound represented by the general formula (17), which is a starting material, can be synthesized in accordance with a known method [for example, a method described in Organic Synthesis Collect, Volume III, 586 (1955); Journal of Chemical Society, 1808 (1951); Synthesis, 793 (1992); Japanese Patent Application Laid-Open No. 325263/1996; or the like]. This compound or a salt thereof is reacted with a halide (iodide, bromide or chloride), whereby the compound (1-5) according to the present invention can be obtained. This reaction is carried out in the presence of a base in a proper solvent. Examples of the solvent used include dimethylformamide, dimethyl sulfoxide, methanol, ethanol, ethoxyethanol, tetrahydrofuran, dioxane and acetonitrile. Examples of the base used include triethylamine, N,N-diisopropylethylamine, pyridine, sodium hydroxide, potassium hydroxide, potassium carbonate, sodium hydrogencarbonate, sodium hydride and potassium hydride. The reaction is performed at room temperature or under heating with stirring.

Step 12:

[0047] The compound (1-5) according to the present invention is hydrolyzed in the presence of proper solvent and base, whereby the compound (1-6) according to the present invention can be obtained as its corresponding carboxylic acid. Alternatively, the compound (1-5) according to the present invention is reacted with any of various amines in the presence or absence of a solvent, or the above carboxylic acid is converted into an active derivative (for example, an acid halide or mixed acid anhydride) and then reacted with any of various amines, whereby the compound (1-6) according to the present invention can be prepared. With respect to the solvent and base used in the hydrolysis, the same conditions as those in Step 6 of Preparation Process 1 may be used.

<Preparation Process 5>


Case of $R^1 = H$ and $X = S$ in the general formula (1):



wherein W , A , R^2 , R^3 and Y^1 have the same meaning as defined above.

[0048] More specifically, a compound represented by a general formula (20) is hydrolyzed under basic conditions and then reacted with its corresponding halide, whereby a compound (1-7) according to the present invention (Step 13). This compound is further converted into an active carboxylic acid derivative and then reacted with any of various amines and alcohols, whereby a compound (1-8) according to the present invention can be prepared (Step 14). The individual steps will hereinafter be described.

<Preparation Process 6> Case of X = O and R¹ = alkyl in the general formula (1):



wherein W, A, A¹, A², Y³, R¹, R², R³ and Q¹ have the same meanings as defined above.

[0052] More specifically, after a compound (21) is reacted with a compound (22) to form a compound (23) (Step 15), exactly the same reactions as in Steps 4, 5, 6 and 7 of Preparation Process 2 are conducted, whereby a compound (24)


can be prepared. After the terminal hydroxy group of the resultant compound (24) is converted into a leaving group (Q^1) in the same manner as in Step 8 of Preparation Process 2, the resultant compound is allowed to react in the same manner as in Step 1 of Preparation Process 2, whereby a compound (1-10) according to the present invention can be prepared. When A^1 in the compound represented by the general formula (24) is a single bond, the compound is subjected to the Mitsunobu reaction with a compound represented by the general formula (3) in the same manner as in Step 9 of Preparation Process 2, whereby a compound (1-9) according to the present invention can be prepared. Step 15 will hereinafter be described in detail.

Step 15:

[0053] The compound (21), which is a starting material, can be purchased as a commercially available reagent or prepared from the compound (9) in accordance with a known method [for example, a method described in Journal of Organic Chemistry, 21, 1149 (1956); Angewandte Chemie, 80, 364 (1968); or the like]. This compound (21) is heated and stirred in the presence of an acid anhydride (22) and a base in accordance with a method described in literature [for example, Journal of American Chemical Society, 72, 1988 (1950); Journal of American Chemical Society, 73, 4911 (1951); Journal of Medicinal Chemistry, 39, 3897 (1996); or the like], whereby the compound (23) can be prepared. Examples of the base used include pyridine, sodium acetate and potassium acetate.

<Preparation Process 7>

Case of $R^1 = \text{group } COY^2$ in the general formula (1):



wherein R^2 , R^3 , Y^1 , Y^2 , A and W have the same meaning as defined above, and X^3 is O, S or NH.

[0054] More specifically, a compound represented by the general formula (5) is condensed with a malonic acid derivative represented by a general formula (26) in the same manner as in Step 10 of Preparation Process 4, thereby obtaining a compound represented by a general formula (27). The compound thus obtained is reduced in the same manner as in Step 7 of Preparation Process 2 to convert it into a compound represented by a general formula (28), and the compound (28) is then subjected to a brominating reaction, whereby a compound represented by a general formula (29) can be prepared (Step 16). Finally, the compound (29) is reacted with any of various nucleophilic reagents, whereby a compound (1-11) according to the present invention can be prepared (Step 17). Steps 16 and 17 will hereinafter be described in detail.


Step 16:

[0055] The compound represented by the general formula (28) is reacted with bromine in the presence of a proper solvent in accordance with a known method [for example, a method described in Organic Synthesis Collect Volume III, 705 (1955); Organic Synthesis Collect Volume I, 245 (1945); Tetrahedron Letters, 24, 163 (1983); or the like], whereby the compound represented by the general formula (29) can be prepared. Examples of the solvent used include acetic acid, diethyl ether, dioxane and carbon tetrachloride.

Step 17:

[0056] The compound represented by the general formula (29) is reacted with any of various alcohols, amines and thiols in the presence of proper solvent and base, whereby the compound (1-11) according to the present invention can be prepared. No particular limitation is imposed on the solvent used so far as it does not affect the reaction. Examples thereof include various kinds of alcohols, dioxane, tetrahydrofuran, dimethylformamide and dimethyl sulfoxide. Examples of the base used include sodium hydroxide, potassium hydroxide, potassium carbonate, sodium hydrogen-carbonate, sodium hydride, triethylamine and pyridine.

<Preparation Process 8>



wherein R^2 , R^3 , X^1 , Y^1 , Y^2 , A^2 , Q^1 and W have the same meanings as defined above.

[0057] More specifically, a compound represented by a general formula (30) is reacted with a compound represented by a general formula (31) in the same manner as in Step 1 of Preparation Process 1, thereby obtaining a compound (32). The compound thus obtained is then catalytically reduced in the same manner as in Step 7 of Preparation Process 2 to obtain a compound (33). Finally, the compound (33) is reacted with the compound (3), whereby a compound (1-12) according to the present invention can be prepared.

[0058] The compound (1) according to the present invention obtained by each of the above-described preparation processes can be isolated in the form of crystals or a liquid, as needed, by the conventional means for isolation and purification, for example, recrystallization, distillation and/or chromatography. The compound may also be converted into a salt or solvate as needed.

[0059] The compounds (1) according to the present invention are excellent in the effect of lowering blood glucose and lipid and hence useful as medicines for preventing or treating diabetes mellitus, hyperlipemia, obesity and the like.

[0060] The medicine according to the present invention is prepared by formulating an effective amount of the compound (1) or the salt thereof as an active ingredient in suitable combination with pharmaceutically acceptable, known carriers, for example, excipients, binders, disintegrators, lubricants, dissolution aids, suspending agents and the like according to pharmacological effects intended, administration object, administration end, administration form, etc. Examples of administration forms include oral administrations by tablets, capsules, granules, powder, syrup, etc., and parenteral administrations by injections, ophthalmic solutions suppositories, etc. In the medicine according to the present invention, the dose of the compound (1) varies according to the condition, age and weight of a patient to be administered, and the administration method thereof. However, the compound (1) may be generally administered in a

dose of 0.1 to 1,000 mg per day for an adult.

[0061] The compounds (1) according to the present invention may also be used as veterinary medicines for other mammals than the human.

[0062] The present invention will hereinafter be described more specifically by the following Preparation Examples, Examples and Test Example. However, these examples are intended to illustrate the present invention and by no means limit the present invention.

Preparation Example 1:

Preparation of 4-[2-(1-oxo-1,2-dihydrophthalazin-2-yl)ethoxy]benzaldehyde (Compound 5):

[0063] After dimethylformamide (100 ml) was added to a mixture of 1-phthalazinone (1.46 g), 4-[2-(methanesulfonyloxy)ethoxy]benzaldehyde (2.44 g) and potassium carbonate (2.07 g), and the mixture was heated and stirred for 4 hours at 85 to 90°C on an oil bath, the reaction mixture was poured into ice water and extracted with ethyl acetate. After the resultant extract was washed with water and then with brine, it was dried over anhydrous sodium sulfate, and the solvent was removed under reduced pressure. The resultant residue was purified by column chromatography on silica gel. A chloroform eluate was crystallized from ether to obtain the title compound (2.17 g, yield: 73.8%) as colorless crystals.

¹H-NMR (CDCl₃, δ):

4.52(2H,t), 4.68(2H,t), 7.00(2H,d), 7.71-7.86(5H,m), 8.19(1H,s), 8.44(1H,dd), 9.86(1H,s).

Preparation Example 2:

Preparation of 4-[2-(4-oxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)ethoxy]benzaldehyde (Compound 5):

[0064] Dimethyl sulfoxide (100 ml) was added to 2-(4-oxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)ethanol (11.6 g) and p-fluorobenzaldehyde (11.2 g) to prepare a solution. After sodium hydride (3.6 g; 60% assay) was added portionwise to the solution under ice cooling, the mixture was stirred at room temperature for 2.5 hours. Thereafter, the reaction mixture was poured into ice water, and the precipitate was collected by filtration, washed with water and dried under reduced pressure. After ether was added to the thus-obtained crude crystals, and the resultant mixture was heated and stirred, the mixture was cooled down to room temperature. The precipitate was collected by filtration and dried to obtain the title compound (15.4 g, yield: 86.3%) as colorless crystals.

¹H-NMR (CDCl₃, δ):

3.99(2H,t), 4.36(2H,t), 5.37(2H,s), 6.97-7.01(3H,m), 7.12(1H,brt), 7.45(1H,ddd), 7.84(2H,d), 7.95(1H,ddd), 9.89(1H,s).

Preparation Example 3:

Preparation of 2-[2-[4-(2-methoxyvinyl)phenoxy]ethyl]-1,2-dihydro-1-phthaladinone (Compound 6):

[0065] (Methoxymethyl)triphenylphosphonium chloride (13.71 g, 40 mmol) and diisopropylamine (4.22 ml, 30 mmol) were suspended in anhydrous tetrahydrofuran (100 ml). A 1.58 M hexane solution (19 ml, 30 mmol) of n-butyllithium was added to the suspension at -10°C, and the mixture was stirred for 1 hour. A solution of 4-[2-(1-oxo-1,2-dihydrophthalazin-2-yl)ethoxy]benzaldehyde (Compound 5) (5.88 g, 20 mmol) in tetrahydrofuran (40 ml) was then added dropwise to the mixture at the same temperature, and the resultant mixture was stirred for 2 hours at -10°C to room temperature. After completion of the reaction, the reaction mixture was poured into ice water and extracted with ethyl acetate. The resultant extract was dried over anhydrous sodium sulfate, and the solvent was removed under reduced pressure. The resultant residue was purified by column chromatography on silica gel (hexane:ethyl acetate = 4:1) to obtain the title compound (a mixture of E/Z) (6.5 g, yield: 99%) as pale yellow crystals.

¹H-NMR (CDCl₃, δ):

3.64(1.5H,s), 3.73(1.5H,s), 4.34-4.70(4H,m), 5.10(0.5H,d), 5.75(0.5H,d), 6.05(0.5H,d), 6.80-7.51(4.5H,m), 7.68-7.84(3H,m), 8.17(1H,s), 8.39-8.48(1H,m).

Preparation Example 4:

Preparation of 2-[2-[4-(2,2-diethoxyethyl)phenoxy]ethyl]-1,2-dihydro-1-phthaladinone (Compound 7):

- 5 [0066] Ethanol (200 ml) was added to 2-[2-[4-(2-methoxyvinyl)phenoxy]ethyl]-1,2-dihydro-1-phthaladinone (Compound 6) (6.44g, 20 mmol) and p-toluenesulfonic acid hydrate (0.38 g, 2 mmol), and the mixture was heated under reflux for 6 hours. After completion of the reaction, the solvent was removed, and ethyl acetate was added to the resultant residue. After the resultant mixture was washed successively with a 5% aqueous solution of sodium hydrogencarbonate and brine and dried over anhydrous magnesium sulfate, the solvent was removed to obtain the title compound
- 10 (7.2 g, yield: 94.2%) as a pale yellow oil.

¹H-NMR (CDCl₃, δ):

1.15(6H,t), 2.85(2H,d), 3.34-3.75(4H,m), 4.32-4.72(5H,m), 6.84(2H,d), 7.10(2H,d), 7.69-7.85(3H,m), 8.18(1H,s), 8.34-8.50(1H,m).

Preparation Example 5:

Preparation of 2-ethoxy-3-[4-[2-(1-oxo-1,2-dihydrophthalazin-2-yl)ethoxy]phenyl]propionitrile (Compound 8):

- 20 [0067] 2-[2-[4-(2,2-Diethoxyethyl)phenoxy]ethyl]-1,2-dihydro-1-phthaladinone (Compound 7) (7.2 g, 18.8 mmol) was dissolved in dichloromethane (100 ml). Trimethylsilyl nitrile (7.52 ml, 56.4 mmol) and boron trifluoride etherate (0.58 ml, 4.7 mmol) were successively added to the solution, and the mixture was stirred for 1 hour. After completion of the reaction, dichloromethane was added to the reaction mixture. After the resultant mixture was washed with a 5% aqueous solution of sodium hydrogencarbonate and water and dried over anhydrous magnesium sulfate, the solvent was
- 25 removed under reduced pressure. The resultant residue was purified by column chromatography on silica gel (hexane:ethyl acetate = 4:1) to obtain the title compound (6.0 g, yield: 88.0%) as colorless crystals.

¹H-NMR (CDCl₃, δ):

1.22(3H,t), 3.05(2H,d), 3.30-4.00(2H,m), 4.19(1H,t), 4.40-4.70(4H,m), 6.85(2H,d), 7.15(2H,d), 7.68-7.84(3H,m), 8.17(1H,s), 8.39-8.50(1H,m).

Preparation Example 6:

Preparation of 2-ethoxy-3-[4-(2-hydroxyethyl)phenyl]propanamide (Compound 11):

- 35 [0068] 2-Ethoxy-3-[4-(2-benzyloxyethyl)phenyl]propanamide (Compound 10) (13.04 g) was dissolved in ethanol (150 ml) and acetic acid (30 ml). To the solution, was added 10% palladium on charcoal (8 g), followed by hydrogenation at room temperature for 4 hours. After completion of the reaction, the reaction mixture was filtered through Celite, and the filtrate was concentrated under reduced pressure. The resultant residue was purified by column chromatography on silica gel (CHCl₃:MeOH = 100:2) to obtain the title compound (8.38 g, yield: 88.6%) as a colorless oil.
- 40

¹H-NMR (CDCl₃, δ):

1.14(3H,t), 2.71-4.04(10H,m), 6.18(1H,brs), 6.52(2H,brs), 7.08-7.32(4H,m).

45 Preparation Example 7:

Preparation of 2-ethoxy-3-[4-[2-(methanesulfonyloxy)ethyl]phenyl]propanamide (Compound 12):

- 50 [0069] 2-Ethoxy-3-[4-(2-hydroxyethyl)phenyl]propanamide (Compound 11) (8.38 g, 35.3 mmol) and triethylamine (4.29 g, 42.4 mmol) were dissolved in methylene chloride (80 ml). Methanesulfonyl chloride (4.85 g, 42.4 mmol) was added dropwise to the solution under ice cooling, and the resultant mixture was stirred at room temperature for 1 hour. After completion of the reaction, the solvent was removed under reduced pressure, and the residue was dissolved in ethyl acetate. The solution was washed with a saturated aqueous solution of sodium hydrogencarbonate, water and brine in that order, and dried over anhydrous magnesium sulfate. The solvent was then removed under reduced pressure to obtain the title compound (8.44 g, yield: 75.8%) as colorless crystals.
- 55

¹H-NMR (CDCl₃, δ):

1.14(3H,t), 2.78-3.64(9H,m), 3.81-4.05(1H,m), 4.41(2H,t), 5.74(1H,brs), 6.48(1H,brs), 7.08-7.23(4H,m).

Preparation Example 8:

Preparation of 3-[4-(benzyloxy)phenyl]-2-methoxy-2-propenenitrile (Compound 14):

- 5 [0070] 4-Benzyloxybenzaldehyde (2.12 g, 10 mmol) and methoxyacetonitrile (748 mg, 10 mmol) were dissolved in dimethylformamide (30 ml). Sodium hydride (60% assay; 480 mg, 12 mmol) was added to the solution at room temperature, and the resultant mixture was heated and stirred for 1 hour at 110°C on an oil bath. After completion of the reaction, the reaction mixture was poured into ice water and extracted with ethyl acetate. After the resultant extract was washed with brine and dried over anhydrous magnesium sulfate, the solvent was removed under reduced pressure.
- 10 The resultant residue was purified by column chromatography on silica gel (hexane:ethyl acetate = 10:1) to obtain the title compound (a mixture of E/Z; 1.58 g, yield: 59.7%) as colorless syrup.

¹H-NMR (CDCl₃, δ):

- 3.76(1.5H,s), 3.90(1.5H,s), 5.09(2H,s), 6.14(0.5H,s), 6.53(0.5H,s), 6.95(1H,d), 6.98(1H,d), 7.10-
 15 7.40(5H,m), 7.51(1H,d), 7.58(1H,d).

Example 1:

- Preparation of 2-ethoxy-3-[4-[2-(1-oxo-1,2-dihydrophthalazin-2-yl)ethoxy]phenyl]propanamide [Invention Compound
 20 (1A)]:

- [0071] 2-Ethoxy-3-[4-[2-(1-oxo-1,2-dihydrophthalazin-2-yl)ethoxy]phenyl]propionitrile (Compound 8) (6.0 g, 16.5 mmol) was dissolved in ethanol (200 ml), and a 6N aqueous solution (8.25 ml, 49.5 mmol) of sodium hydroxide was added to the solution. The resultant mixture was heated under reflux for 1.5 hours. After completion of the reaction,
 25 water was added to the reaction mixture to conduct extraction with ethyl acetate. The resultant extract was washed with brine and dried over anhydrous magnesium sulfate. The solvent was then removed under reduced pressure. The resultant residue was purified by column chromatography on silica gel (hexane:ethyl acetate = 4:1) and recrystallized from ethyl acetate to obtain Invention Compound (1A) (3.3 g, yield: 52.5%) as colorless crystals.

- 30 Example 2:

[0072] Invention Compounds (1D), (1F), (1H), (1J), (1K), (1N), (1AB), (1AC), (1BG), (1BU) and (1BV) were prepared in the same manner as in Example 1.

- 35 Example 3:

Preparation of 2-ethoxy-3-[4-[2-(4-oxo-3,4-dihydroquinazolin-3-yl)ethyl]phenyl]propanamide [Invention Compound
 (1Q)]:

- 40 [0073] 4-Quinazoline (500 mg, 3.42 mmol), 2-ethoxy-3-[4-[2-(methanesulfonyloxy)ethyl]phenyl]propanamide (Compound 12) (1.13 g, 3.42 mmol) and potassium carbonate (1.42 g, 10.3 mmol) were dissolved in dimethylformamide (20 ml), and the resultant solution was heated and stirred at 80°C for 2 hours. After cooling the reaction mixture, water was added to the reaction mixture to conduct extraction with ethyl acetate. The resultant extract was washed with water and then dried over anhydrous magnesium sulfate. The solvent was removed under reduced pressure. The precipitate was
 45 collected by filtration and washed with ether. The crude crystals thus obtained were recrystallized from ethanol to obtain Invention Compound (1Q) (1.11 g, yield: 87.6%).

Example 4:

- 50 [0074] Invention Compounds (1E), (1G), (1I), (1S), (1T), (1U), (1V), (1W), (1X), (1Y), (1AF), (1AI) and (1BR) were prepared in the same manner as in Example 3.

Example 5:

- 55 Preparation of 2-methoxy-3-[4-[2-(4-oxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)ethoxy]phenyl]propanamide [Invention Compound (1P)]:

[0075] 2-(4-Oxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)ethanol (794 mg, 4.11 mmol), 2-methoxy-3-(4-hydroxyphenyl)-

propanamide (802 mg, 4.11 mmol) and triphenylphosphine (1.13 g, 4.31 mmol) were dissolved in tetrahydrofuran (20 ml), and a 40% toluene solution of diethyl azodicarboxylate (1.88 g, 4.31 mmol) was added dropwise to the solution under ice cooling. The resultant mixture was stirred overnight at room temperature. After completion of the reaction, the solvent was removed under reduced pressure. Ether was added to the residue, and the resultant mixture was washed with a 10% aqueous solution of sodium hydroxide and brine. An organic layer was dried over anhydrous magnesium sulfate. The solvent was then removed under reduced pressure, and the resultant residue was purified by column chromatography on silica gel (hexane:ethyl acetate = 4:1) to obtain Invention Compound (1P) (976 mg, yield: 64.1%) as colorless crystals.

10 Example 6:

[0076] Invention Compounds (1R), (1AH), (1AJ), (1AK), (1AL), (1AM), (1AS), (1AT), (1AV), (1AY), (1AZ) and (1BE) were prepared in the same manner as in Example 5.

15 Example 7:

Preparation of ethyl 2-ethylamino-3-{4-[2-(4-oxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)ethoxy]phenyl}propionate [Invention Compound (1L)]:

20 [0077] Ethyl 2-amino-3-{4-[2-(4-oxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)ethoxy]phenyl}propionate hydrochloride (1.05 g, 2.5 mmol) was dissolved in ethanol (10 ml), and ethyl iodide (390 mg, 2.5 mmol) and diisopropylethylamine (323 mg, 2.5 mmol) were added dropwise to the solution. The resultant mixture was stirred for 2 days at 60°C [during which additional ethyl iodide and diisopropylethylamine (each, 5 mmol) were added]. After completion of the reaction, the solvent was removed, and ethyl acetate was added to the residue. The resultant mixture was washed with water and dried over anhydrous magnesium sulfate. The solvent was then removed under reduced pressure, and the resultant residue was purified by column chromatography on silica gel to obtain Invention Compound (1L) (627 mg, yield: 60.8%) as a colorless oil.

30 Example 8:

Preparation of 2-ethylamino-3-{4-[2-(4-oxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)ethoxy]phenyl}propionic acid [Invention Compound (1M)]:

35 [0078] A 1N aqueous solution (5 ml) of sodium hydroxide was added to a solution of ethyl 2-ethylamino-3-{4-[2-(4-oxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)ethoxy]phenyl}propionate (627 mg) in ethanol (8 ml), and the resultant mixture was stirred overnight at 60°C. After the completion of a reaction was confirmed by using TLC, the reaction mixture was concentrated under reduced pressure. Water was added to the residue, and the resultant mixture was neutralized with 3% hydrochloric acid. The precipitate was then collected by filtration, washed with water and then dried to obtain Invention Compound (1M) (640 mg, quantitative) in the form of the hydrochloride as colorless crystals.

40 Example 9:

[0079] Invention Compounds (1BA), (1BB) and (1BF) were prepared in the same manner as in Example 8.

45 Example 10:

Preparation of 2-ethylthio-3-{4-[2-(4-oxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)ethoxy]phenyl}propanoic acid [Invention Compound (1O)]:

50 [0080] A 15% aqueous solution (60 ml) of sodium hydroxide was added to 5-{4-[2-(4-oxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)ethoxy]benzyl}-2,4-thiazolidinedione (3.0 g). After the resultant mixture was gently heated to reflux for 30 minutes, methanol (60 ml) was then added to the mixture at room temperature, and a methanol solution of ethyl iodide (5.3 g) was then added dropwise, followed by stirring for 1.5 hours. Thereafter, the reaction mixture was poured into ice water, acidified with hydrochloric acid and extracted with ethyl acetate. The resultant extract was washed with brine and dried over anhydrous sodium sulfate. The solvent was then removed under reduced pressure. The resultant residue was purified by column chromatography on silica gel (chloroform:methanol = 50:1) and crystallized from hexane-ethyl acetate to obtain Invention Compound (1O) (600 mg, yield: 19.9%) as colorless crystals.

Example 11:

Preparation of 2-ethoxy-3-[4-[2-(1-oxo-1,2-dihydrophthalazin-2-yl)ethoxy]phenyl]propanoic acid [Invention Compound (1B)]:

[0081] 2-Ethoxy-3-[4-[2-(1-oxo-1,2-dihydrophthalazin-2-yl)ethoxy]phenyl]propanamide [Invention Compound (1A)] (12.55 g, 32.9 mmol) was dissolved in dioxane (148 ml), and 2N hydrochloric acid (16.5 ml, 32.9 mmol) and a catalytic amount of TiCl_4 were added to the solution, followed by stirring at 110°C for 6 hours. After completion of the reaction, the solvent was removed, and chloroform was added to the residue. The precipitate was removed by filtration, and the filtrate was dried over anhydrous magnesium sulfate. The solvent was then removed, and the resultant residue was purified by column chromatography on silica gel (hexane:ethyl acetate = 2:3) and recrystallized from ethyl acetate to obtain Invention Compound (1B) (5.23 g, yield: 41.8%) as colorless crystals.

Example 12:

[0082] Invention Compounds (1AE), (1BC), (1BD), (1BJ), (1BK), (1BL) and (1BS) were prepared in the same manner as in Example 11.

Example 13:

Preparation of ethyl 2-ethoxy-3-[4-[2-(1-oxo-1,2-dihydrophthalazin-2-yl)ethoxy]phenyl]propanoate [Invention Compound (1C)]:

[0083] 2-Ethoxy-3-[4-[2-(1-oxo-1,2-dihydrophthalazin-2-yl)ethoxy]phenyl]propanamide [Invention Compound (1A)] (1.0 g, 2.62 mmol) was dissolved in ethanol (30 ml), and 1N hydrochloric acid (2.6 ml, 2.6 mmol) and a catalytic amount of TiCl_4 were added to the solution, followed by refluxing for 5 hours. After completion of the reaction, the solvent was removed, and chloroform was added to the residue. The precipitate was removed by filtration, and the filtrate was dried over anhydrous magnesium sulfate. The resultant residue was then purified by column chromatography on silica gel (hexane:ethyl acetate = 2:3) to obtain Invention Compound (1C) (0.2 g, yield: 20.0%) as a colorless oil.

Example 14:

[0084] Invention Compounds (1AN), (1AO), (1AP), (1AQ), (1AU), (1AW), (1AX) and (1BO) were prepared in the same manner as in Example 13.

Example 15:

Preparation of 2-ethoxy-3-[4-[2-(1-oxo-1,2-dihydrophthalazin-2-yl)ethoxy]phenyl]propanohydroxamic acid [Invention Compound (1AA)]:

[0085] 2-Ethoxy-3-[4-[2-(1-oxo-1,2-dihydrophthalazin-2-yl)ethoxy]phenyl]propanoic acid [Invention Compound (1B)] (500 mg, 1.31 mmol) was dissolved in dimethylformamide (10 ml), and carbonyldiimidazole (233 mg, 1.44 mmol) was added to the solution. After the resultant mixture was stirred at room temperature for 1 hour, hydroxylamine hydrochloride (200 mg, 2.88 mmol) was added, and the mixture was stirred at the same temperature for 12 hours. After completion of the reaction, water was added to the reaction mixture to conduct extraction with ethyl acetate. The resultant extract was washed with water and then with brine, and dried over anhydrous magnesium sulfate. The solvent was then removed, and the residue was purified by column chromatography on silica gel (ethyl acetate:hexane = 3:2 to 4:1) and recrystallized from a mixed solvent of ethyl acetate and ether to obtain Invention Compound (1AA) (150 mg, yield: 28.8%) as colorless crystals.

Example 16:

[0086] Invention Compounds (1Z), (1AD), (1AG), (1BH) and (1BP) were prepared in the same manner as in Example 15.

Example 17:

Preparation of ethyl 2-ethoxy-3-{4-[2-(4-methyl-1-oxo-1,2-dihydrophthalazin-2-yl)ethoxy]phenyl}propanoate [Invention Compound (1BI)]:

[0087] Ethyl 2-ethoxy-3-{4-[2-(4-methyl-1-oxo-1,2-dihydrophthalazin-2-yl)ethoxy]phenyl}propanoate (4.96 g, 11.75 mmol) was dissolved in ethyl acetate (60 ml), and 10% palladium on carbon (600 mg) was added to conduct catalytic reduction for 10 hours. After completion of the reaction, the catalyst was removed by filtration, and the filtrate was concentrated. The resultant residue was purified by column chromatography on silica gel to obtain Invention Compound (1BI) (4.54 g, yield: 91.2%) as colorless crystals.

Example 18:

[0088] Invention Compounds (1AR), (1BM), (1BN), (1BQ), (1BT) and (1BW) were prepared in the same manner as in Example 17.

[0089] The structures and physical data of the compounds obtained in the above-described Examples are shown in the following tables. W, A, R¹, R², R³, X and Y¹ in the tables mean the respective symbols in the general formula (1). The position of substitution (abbreviated as "PS" in the tables) indicates the position of W-A- substituted on a benzene ring.

Table 2






No.	W	A	PS	R ¹	R ²	R ³	X	Y ¹	m. p. (°C)	NMR (CDC l ₃ , δ)
1E		-Cl-CH ₂ -O-	4	II	Et	II	O	NH ₂	141-145	1.12 (3H, t), 2.60 (3H, s), 2.85 (1H, dd), 3.06 (1H, dd), 3.36-3.53 (2H, m), 3.86 (1H, dd), 4.39 (2H, t), 4.60 (2H, t), 5.42 (1H, br), 6.42 (1H, br), 6.84 (2H, d), 7.12 (2H, d), 7.74-7.84 (3H, m), 8.46-8.48 (1H, m)
1F		-Cl-CH ₂ -O-	4	II	Et	II	O	NH ₂	87-88	1.11 (3H, t), 2.83 (1H, dd), 3.05 (1H, dd), 3.32-3.52 (2H, m), 3.85 (1H, dd), 4.44 (2H, t), 4.71 (2H, t), 5.59 (1H, brs), 6.42 (1H, brs), 6.85 (2H, d), 7.12 (2H, d), 7.52-7.59 (5H, m), 7.72-7.81 (3H, m), 8.53-8.55 (1H, m)
1G		-Cl-CH ₂ -O-	4	II	Et	II	O	NH ₂	138-140	1.11 (3H, t), 2.83 (1H, dd), 3.05 (1H, dd), 3.35-3.52 (2H, m), 3.85 (1H, dd), 4.45 (2H, t), 4.73 (2H, t), 5.51 (1H, br), 6.41 (1H, br), 6.84 (2H, d), 7.12 (2H, d), 7.40-7.43 (1H, m), 7.76-7.90 (4H, m), 8.42-8.45 (1H, m), 8.50-8.53 (1H, m), 8.75-8.77 (1H, m)
1H		 -Cl-CH ₂ -O-	4	II	Et	II	O	NH ₂	104-106	1.14 (3H, t), 2.86 (1H, dd), 3.08 (1H, dd), 3.40-3.55 (2H, m), 3.80 (1H, d), 3.89 (1H, dd), 4.05-4.13 (2H, m), 4.47-4.62 (3H, m), 5.40 (1H, br), 6.43 (1H, br), 6.85 (2H, d), 7.15 (2H, d), 7.72-7.74 (1H, m), 7.79-7.87 (2H, m), 8.21 (1H, s), 8.44 (1H, d)

Table 3





No.	W	A	PS	R ¹	R ²	R ³	X	Y ¹	m.p. (°C)	NMR (CDCl ₃ , δ)
1I		-CH ₂ -CH ₂ -	4	H	Et	H	O	NH ₂	146-149	1.13(3H, t), 2.86(1H, dd), 3.09-3.15(3H, m), 3.40-3.53(2H, m), 3.90(1H, dd), 4.43-4.47(2H, m), 5.53(1H, br), 6.44(1H, br), 7.18(2H, d), 7.20(2H, d), 7.68-7.83(3H, m), 8.16(1H, s), 8.42(1H, dd)
1J		-CH ₂ -CH ₂ -O-	4	H	Et	H	O	NH ₂	101-102	1.13(3H, t), 2.86(1H, dd), 3.07(1H, dd), 3.38-3.54(2H, m), 3.87(1H, dd), 3.95(2H, t), 4.18(2H, t), 5.3-5.3 (1H, brs), 5.37(2H, s), 6.40(1H, brs), 6.79(2H, d), 6.97(1H, dd), 7.09-7.14(1H, m), 7.15(2H, d), 7.42-7.46(1H, m), 7.95(1H, dd)
1K		-CH ₂ -CH ₂ -O-	4	H	Et	H	O	OH	98-99	1.16(3H, t), 2.94(1H, dd), 3.06(1H, dd), 3.38-3.46(1H, m), 3.55-3.63(1H, m), 3.95(2H, t), 4.02(1H, dd), 4.17(2H, t), 5.37(2H, s), 6.80(2H, d), 6.97(1H, d), 7.10(1H, dt), 7.42-7.46(1H, m), 7.95(1H, dd)
1L		-CH ₂ -CH ₂ -O-	4	H	Et	H	NH	OH	011	1.06(3H, t), 1.15(3H, t), 2.48-2.58(1H, m), 2.60-2.67(1H, m), 2.82-2.93(2H, m), 3.46(1H, t), 3.94(2H, t), 4.09(2H, q), 4.17(2H, t), 5.36(2H, s), 6.79(2H, d), 6.96(1H, d), 7.07-7.12(3H, m), 7.43(1H, dt), 7.95(1H, dd)

Table 4






No	W	A	PS	R ¹	R ²	R ³	X	Y ¹	m.p. (°C)	NMR (CDCl ₃ , δ)
IM		-CH ₂ -CH ₂ -O-	4	H	H ¹	H	NH	OH	228-229 (HCl salt)	0.99 (3H, t), 2.38-2.49 (1H, m), 2.50-2.55 (1H, m), 2.72-2.83 (2H, m), 3.26 (1H, t), 3.72 (2H, t), 4.03 (2H, t), 4.81 (2H, s), 6.74-6.77 (3H, m), 6.97 (1H, t), 7.09 (2H, d), 7.30 (1H, dt), 7.66 (1H, dd), (D ₂ O+NaOD) (HCl salt)
IN		-CH ₂ -CH ₂ -O-	4	H		H	O	NH ₂	82-83	0.95 (3H, d), 1.11 (3H, d), 2.80 (1H, dd), 3.05 (1H, dd), 3.45 (1H, dd), 3.92 (1H, dd), 3.95 (2H, t), 4.18 (2H, t), 5.37 (2H, t), 5.43 (1H, br), 6.49 (1H, br), 6.79 (2H, d), 6.97 (1H, d), 7.11 (1H, dt), 7.16 (2H, d), 7.42-7.46 (1H, m), 7.95 (1H, dd)
IO		-CH ₂ -CH ₂ -O-	4	H	H ¹	H	S	OH	71-72	1.22 (3H, t), 2.61-2.68 (2H, m), 2.89 (1H, dd), 3.13 (1H, dd), 3.47 (1H, dd), 3.94 (2H, t), 4.14 (2H, t), 5.35 (2H, s), 6.79 (2H, d), 6.96 (1H, d), 7.08-7.14 (3H, m), 7.41-7.45 (1H, m), 7.93 (1H, dd)
IP		-CH ₂ -CH ₂ -O-	4	H	Me	H	O	NH ₂	142-143	2.89 (1H, dd), 3.08 (1H, dd), 3.34 (2H, s), 3.81 (1H, dd), 3.95 (2H, t), 4.18 (2H, t), 5.32 (1H, br d), 5.37 (2H, s), 6.32 (1H, br d), 6.80 (2H, d), 6.97 (1H, dd), 7.11 (1H, dt), 7.15 (2H, d), 7.42-7.46 (1H, m), 7.95 (1H, dd)

Table 5





No.	W	A	PS	R ¹	R ²	R ³	X	Y ¹	m. p. (°C)	NMR (CDCl ₃ , δ)
10		-CH ₂ -CH ₂ -O-	4	H	Et	H	O	NH ₂	156-157	1.02(3H, t), 2.70(1H, dd), 2.84(1H, dd), 3.22-3.30(1H, m), 3.39-3.47(1H, m), 3.72(1H, d), 4.26(2H, t), 4.37(2H, t), 6.81-6.85(2H, m), 7.04-7.11(4H, m), 7.55(1H, ddd), 7.69(1H, dd), 7.83(1H, ddd), 8.17(1H, dd), 8.38(1H, s) (DMSO-d ₆)
1R		-CH ₂ -CH ₂ -O-	4	H	Et	H	O	NH ₂	amorphous	1.12(3H, t), 2.84(1H, dd), 3.06(1H, dd), 3.36-3.44(1H, m), 3.46-3.53(1H, m), 3.86(1H, dd), 3.90(2H, t), 4.15(2H, t), 5.18(2H, s), 5.45(1H, brs), 6.42(1H, brs), 6.61-6.64(2H, m), 6.90(1H, dd), 7.00(1H, ddd), 7.09-7.11(2H, m), 7.14-7.18(3H, m), 7.29(1H, ddd), 7.32(2H, ddd), 8.02(1H, dd)
1S		-CH ₂ -CH ₂ -	4	H	Et	H	O	NH ₂	144-145	1.01(3H, t), 2.74(1H, dd), 2.91(1H, dd), 2.99(2H, t), 3.21-3.29(1H, m), 3.40-3.47(1H, m), 3.77(1H, dd), 4.20(2H, t), 7.07-7.16(6H, m), 7.54(1H, ddd), 7.63(1H, dd), 7.81(1H, ddd), 8.13(1H, s), 8.18(1H, dd) (DMSO-d ₆)
1T		-CH ₂ -CH ₂ -O-	4	H	Et	H	O	NH ₂	134-135	1.03(3H, t), 2.71(1H, dd), 2.85(1H, dd), 3.21-3.31(1H, m), 3.41-3.48(1H, m), 3.53(3H, s), 3.74(1H, dd), 4.16(2H, t), 4.33(2H, t), 6.81-6.85(2H, m), 7.05-7.11(4H, m), 7.31(1H, dd), 7.78(1H, ddd), 8.07(1H, dd) (DMSO-d ₆)

Table 6






No.	W	A	PS	R ¹	R ²	R ³	X	Y ¹	m. p. (°C)	NMR (CDCl ₃ , δ)
1U			4	H	Et	H	O	NH ₂	183-184	1.03 (3H, t), 2.71 (1H, dd), 2.85 (1H, dd), 3.22-3.32 (1H, m), 3.41-3.48 (1H, m), 3.55 (3H, s), 3.74 (1H, dd), 3.83 (3H, s), 3.95 (3H, s), 4.15 (2H, t), 4.31 (2H, t), 6.82-6.85 (2H, m), 6.89 (1H, s), 7.05-7.11 (4H, m), 7.44 (1H, s) (DMSO-d ₆)
1V		-CH ₂ -CH ₂ -	4	H	Et	H	O	NH ₂	187-188	1.03 (3H, t), 2.76 (1H, dd), 2.83-2.87 (2H, m), 2.91 (1H, dd), 3.23-3.31 (1H, m), 3.41-3.49 (1H, m), 3.53 (3H, s), 3.78 (1H, dd), 4.12-4.16 (2H, m), 7.08-7.19 (6H, m), 7.45 (1H, dd), 7.77 (1H, ddd), 8.05 (1H, dd) (DMSO-d ₆)
1W			4	H	Et	H	O	NH ₂	112-113	1.03 (3H, t), 1.90 (2H, quin), 2.61 (2H, t), 2.73 (1H, dd), 2.87 (1H, dd), 3.25-3.50 (5H, m), 3.77 (1H, dd), 3.99 (2H, t), 7.06-7.14 (6H, m), 7.29 (1H, ddd), 7.42 (1H, dd), 7.76 (1H, dd), 8.04 (1H, dd) (DMSO-d ₆)

Table 7




No	W	A	PS	R ¹	R ²	R ³	X	Y ¹	m. p. (°C)	NMR (CDCl ₃ , δ)
1X		-CH ₂ -CH ₂ -O-	4	H	Bt	H	O	NH ₂	131-132	1.06(3H, t), 1.14(3H, t), 1.97(3H, s), 2.27(2H, q), 2.84-2.90(1H, m), 3.05-3.09(1H, m), 3.39-3.55(2H, m), 3.81(2H, t), 3.91(1H, dd), 3.94(2H, s), 4.10(2H, t), 5.52(1H, br), 6.45(1H, br), 6.81(2H, d), 7.16(2H, d)
1Y		CH ₂ -CH ₂ -O-	4	H	Bt	H	O	NH ₂	oil	1.14(3H, t), 2.58-2.69(6H, m), 2.85(1H, dd), 3.10(1H, dd), 3.43-3.61(6H, m), 3.75(2H, t), 3.90(1H, dd), 4.08(2H, t), 5.45(1H, br), 6.45(1H, br), 6.78(2H, d), 7.17(2H, d), 7.26-7.34(5H, m)
1Z		-CH ₂ -CH ₂ -O-	4	H	Bt	H	O	NHMe	103-104	1.10(3H, t), 2.75(3H, d), 2.81(1H, dd), 3.06(1H, dd), 3.33-3.48(2H, m), 4.12(1H, dd), 4.40(2H, t), 4.64(2H, t), 6.44(1H, m), 6.83(2H, d), 7.08(2H, d), 7.69-7.84(3H, m), 8.18(1H, s), 8.43-8.45(1H, m)

Table 8






No.	W	A	PS	R ¹	R ²	R ³	X	Y ¹	m.p. (°C)	NMR (CDC l ₃ , δ)
1AA		-CH ₂ -CH ₂ -O-	4	H	Et	H	O	NHOH	145-148	1.09(3H, t), 2.83(1H, dd), 3.07(1H, dd), 3.34-3.49(2H, m), 4.00(1H, dd), 4.40(2H, t), 4.63(2H, t), 6.83(2H, d), 7.08(2H, d), 7.69-7.84(4H, m), 8.19(1H, s), 8.42-8.44(1H, m), 8.85(1H, s)
1AB		-CH ₂ -CH ₂ -O-	4	H	Et	3-OMe	O	NH ₂	160-161	1.13(3H, t), 2.84(1H, dd), 3.07(1H, dd), 3.39-3.53(2H, m), 3.80(3H, s), 3.87(1H, dd), 3.97(2H, t), 4.20(2H, t), 5.43(2H, s), 5.51(1H, br s), 6.45(1H, br s), 6.74-6.80(3H, m), 6.97(1H, dd), 7.10(1H, dt), 7.41-7.46(1H, m), 7.95(1H, dd)
1AC		-CH ₂ -CH ₂ -O-	4	H	Et	3-OMe	O	OH (Na salt)	82-83 (Na salt)	0.98(3H, t), 2.60(1H, dd), 2.84(1H, dd), 3.09-3.18(1H, m), 3.52-3.61(1H, m), 3.70(3H, s), 3.84(2H, t), 4.08(2H, t), 5.42(2H, s), 6.69(1H, dd), 6.82-6.86(2H, m), 7.05(1H, d), 7.13-7.16(1H, m), 7.52(1H, dt), 7.81(1H, dd), (Na salt DMSO-d ₆)
1AD		-CH ₂ -CH ₂ -O-	4	H	Et	H	O		oil	1.17(3H, t), 1.66-1.76(4H, m), 2.94-2.99(3H, m), 3.31-3.54(5H, m), 4.11(1H, t), 4.39(2H, t), 4.64(2H, t), 6.83(2H, d), 7.12(2H, d), 7.69-7.84(3H, m), 8.18(1H, s), 8.43-8.45(1H, m)

Table 9




No.	W	A	PS	R ¹	R ²	R ³	X	Y ¹	m. p. (°C)	NMR (CDCl ₃ , δ)
1AE		-CH ₂ -CH ₂ -O-	4	H	Et	H	O	OH	89-90 -	1.16(3H, t), 2.91(1H, dd), 3.05(1H, dd), 3.38-3.46(1H, m), 3.55-3.63(4H, m), 4.02(1H, dd), 4.25(2H, t), 4.53(2H, t), 7.10-7.14(2H, m), 7.20-7.29(2H, m), 7.69(1H, ddd), 8.24(1H, dd)
1AF		-CH ₂ -CH ₂ -O-	4	H	Et	H	O	NH ₂	172-173	1.03(3H, t), 2.72(1H, dd), 2.86(1H, dd), 3.24-3.31(1H, m), 3.41-3.49(1H, m), 3.74(1H, dd), 4.20(2H, t), 4.26(2H, t), 6.82(2H, d), 7.06(1H, br s), 7.10-7.12(3H, m), 7.41-7.46(2H, m), 7.80-7.84(1H, m), 8.00(1H, dd) (DMSO-d ₆)
1AG		-CH ₂ -CH ₂ -O-	4	H	Et	H	O	NH ₂ OH	146-148	1.09(3H, t), 2.82(1H, dd), 3.07(1H, dd), 3.34-3.38(1H, m), 3.43-3.49(1H, m), 3.61(3H, s), 3.98(1H, dd), 4.25(2H, t), 4.53(2H, t), 6.82-6.85(2H, m), 7.08-7.10(2H, m), 7.18-7.29(2H, m), 7.69(1H, ddd), 8.23(1H, dd), 8.34(1H, br), 8.90(1H, br s)

Table 10





No.	W	A	PS	R ¹	R ²	R ³	X	Y ¹	m. p. (°C)	NMR (CDCl ₃ , δ)
IAH		-CH ₂ -CH ₂ -O-	4	H	CH ₂ CH ₃	H	O	NH ₂	155-156.5	1.05 (3H, t), 2.74 (1H, dd), 2.88 (1H, dd), 3.21-3.37 (3H, m), 3.42-3.50 (1H, m), 3.59 (3H, s), 3.76 (1H, dd), 4.47 (2H, t), 6.87-6.91 (2H, m), 7.06 (1H, brs), 7.13-7.15 (3H, m), 7.48 (1H, ddd), 7.59 (1H, dd), 7.78 (1H, ddd), 8.12 (1H, dd) (DMSO-d ₆)
IAI		-CH ₂ -CH ₂ -O-	4	H	CONH ₂	H	CH ₂	OEt	165-168	1.07 (3H, t), 2.16 (1H, dd), 2.35 (1H, dd), 2.65-2.76 (2H, m), 2.91-2.94 (1H, m), 3.53 (3H, s), 3.96 (2H, q), 4.16 (2H, t), 4.33 (2H, t), 6.69 (1H, brs), 6.82-6.85 (2H, m), 7.01-7.04 (2H, m), 7.23 (1H, brs), 7.31 (1H, ddd), 7.45 (1H, dd), 7.78 (1H, ddd), 8.07 (1H, dd) (DMSO-d ₆)
IAJ		-CH ₂ -CH ₂ -O-	4	H	CH ₂ CONH ₂	H	O	OEt	142-144	1.29 (3H, t), 2.89 (1H, dd), 3.10 (1H, dd), 3.61 (3H, s), 3.79 (1H, d), 4.03 (1H, d), 4.20-4.29 (2H, m), 4.32 (2H, t), 4.52 (2H, t), 5.68 (1H, brs), 6.04 (1H, brs), 6.87-6.90 (2H, m), 7.14-7.18 (2H, m), 7.22 (1H, dd), 7.27 (1H, ddd), 7.70 (1H, ddd), 8.22 (1H, dd)
IAK		-CH ₂ -CH ₂ -O-	4	H	CH ₂ CONH ₂	H	O	NH ₂	220-222	2.80 (1H, dd), 2.96 (1H, dd), 3.53 (3H, s), 3.70-3.80 (2H, m), 3.90 (1H, dd), 4.16 (2H, t), 4.33 (2H, t), 6.82-6.86 (2H, m), 6.90-7.18 (5H, m), 7.31 (1H, ddd), 7.46 (1H, dd), 7.52 (1H, brs), 7.79 (1H, ddd), 8.07 (1H, dd) (DMSO-d ₆)

Table 11





No.	W	A	PS	R ¹	R ²	R ³	X	Y ¹	m.p. (°C)	NMR (CDC l ₃ , δ)
1AL		-CH ₂ -CH ₂ -O-	4	H	CH ₂ COOEt	H	O	NH ₂	138-139.5	1.17 (3H, t), 2.81 (1H, dd), 2.93 (1H, dd), 3.53 (3H, s), 3.97 (1H, dd), 3.98 (1H, d), 4.08 (2H, q), 4.09 (1H, d), 4.16 (2H, t), 4.33 (2H, t), 6.81-6.85 (2H, m), 7.11-7.22 (4H, m), 7.31 (1H, ddd), 7.45 (1H, dd), 7.78 (1H, ddd), 8.07 (1H, dd) (DMSO-d ₆)
1AM		-CH ₂ -CH ₂ -O-	4	H	CH ₂ CH ₂ OH	H	O	NH ₂	144-146	2.74 (1H, dd), 2.91 (1H, dd), 3.35-3.50 (4H, m), 3.53 (3H, s), 3.81 (1H, dd), 4.16 (2H, t), 4.33 (2H, t), 4.62 (1H, t), 6.81-6.85 (2H, m), 7.10-7.13 (3H, m), 7.24 (1H, brs), 7.31 (1H, ddd), 7.45 (1H, dd), 7.78 (1H, ddd), 8.07 (1H, dd) (DMSO-d ₆)
1AN		-CH ₂ -CH ₂ -O-	4	H	COOMe	H	CH ₂	OMe	90-92	2.39 (1H, dd), 2.52 (1H, dd), 2.69 (1H, dd), 2.81 (1H, dd), 2.93-2.98 (1H, m), 3.53 (3H, s), 3.54 (3H, s), 3.55 (3H, s), 4.17 (2H, t), 4.33 (2H, t), 6.83-6.86 (2H, m), 7.02-7.05 (2H, m), 7.31 (1H, ddd), 7.45 (1H, dd), 7.78 (1H, ddd), 8.07 (1H, dd) (DMSO-d ₆)
1AO		-CH ₂ -CH ₂ -O-	4	H	CH ₂ COOMe	H	O	OMe	102-103.5	2.87-2.97 (2H, m), 3.53 (3H, s), 3.59 (3H, s), 3.61 (3H, s), 4.08 (1H, d), 4.16 (2H, t), 4.19 (1H, d), 4.26 (1H, t), 4.33 (2H, t), 6.82-6.85 (2H, m), 7.08-7.11 (2H, m), 7.31 (1H, ddd), 7.45 (1H, dd), 7.78 (1H, ddd), 8.07 (1H, dd) (DMSO-d ₆)

Table 12





No.	W	A	PS	R ¹	R ²	R ³	X	Y ¹	m. p. (°C)	NMR (CDC l ₃ , δ)
IAP		-CH ₂ -CH ₂ -O-	4	H	COOEt	H	CH ₂	OEt	oil	1.09 (3H, t), 1.14 (3H, t), 2.37 (1H, dd), 2.49 (1H, dd), 2.69 (1H, dd), 2.80 (1H, dd), 2.91-2.95 (1H, m), 3.53 (3H, s), 2.97-4.04 (4H, m), 4.17 (2H, t), 4.33 (2H, t), 6.83-6.87 (2H, m), 7.03-7.06 (2H, m), 7.31 (1H, ddd), 7.45 (1H, dd), 7.78 (1H, ddd), 8.07 (1H, dd) (DMSO-d ₆)
IAQ		-CH ₂ -CH ₂ -O-	4	H	CH ₂ COOEt	H	O	OEt	oil	1.11 (3H, t), 1.17 (3H, t), 2.92 (2H, d), 3.53 (3H, s), 4.02-4.25 (9H, m), 4.33 (2H, t), 6.82-6.85 (2H, m), 7.09-7.12 (2H, m), 7.31 (1H, ddd), 7.45 (1H, dd), 7.78 (1H, ddd), 8.07 (1H, dd)
IAR		-CH ₂ -CH ₂ -O-	4	H	CH ₂ CH ₃	H	O	OH	119-121	1.16 (3H, t), 2.92 (1H, dd), 3.06 (1H, dd), 3.40-3.48 (1H, m), 3.52-3.65 (1H, m), 3.65 (3H, s), 4.03 (1H, dd), 4.26 (2H, t), 4.54 (2H, t), 6.81-6.85 (2H, m), 7.09-7.14 (2H, m), 7.43 (1H, s), 7.51 (1H, d), 8.37 (1H, d)
IAS		-CH ₂ -CH ₂ -O-	4	H	CH ₂ COOEt	H	O	Nil ₂	116-118	1.16 (3H, t), 2.55 (3H, s), 2.80 (1H, dd), 2.93 (1H, dd), 3.97 (1H, dd), 3.98 (1H, d), 4.04 (2H, q), 4.10 (1H, d), 4.34 (2H, t), 4.36 (2H, t), 6.80-6.84 (2H, m), 7.09-7.14 (3H, m), 7.22 (1H, brs), 7.84-7.95 (3H, m), 8.30 (1H, d) (DMSO-d ₆)

Table 13






No.	W	A	PS	R ¹	R ²	R ³	X	Y ¹	m. p. (°C)	NMR (CDC l ₃ , δ)
1AT		-CH ₂ -CH ₂ -O-	4	H	CH ₂ CH ₂ OH	H	O	Nil ₂	127-128.5	2.56(3H, s), 2.73(1H, dd), 2.90(1H, dd), 3.35-3.47(4H, m), 3.80(1H, dd), 4.34(2H, t), 4.46(2H, t), 4.62(1H, t), 6.80-6.83(2H, m), 7.09-7.12(3H, m), 7.24(1H, brs), 7.85-7.95(3H, m), 8.30(1H, d) (DMSO-d ₆)
1AU		-CH ₂ -CH ₂ -O-	4	H	CH ₂ COOMe	H	O	OMe	oil	2.55(3H, s), 2.90-2.92(2H, m), 3.58(3H, s), 3.61(3H, s), 4.08(1H, d), 4.19(1H, d), 4.25(1H, t), 4.34(2H, t), 4.46(2H, t), 6.81-6.84(2H, m), 7.07-7.10(2H, m), 7.84-7.95(3H, m), 8.28-8.31(1H, m) (DMSO-d ₆)
1AV		-CH ₂ -CH ₂ -O-	4	H	CH ₂ COOEt	H	O	OEt	oil	1.10(3H, t), 1.16(3H, t), 2.55(3H, t), 2.91(2H, d), 4.01-4.11(5H, m), 4.17(1H, d), 4.22(1H, t), 4.34(2H, t), 4.46(2H, t), 6.81-6.84(2H, m), 7.07-7.11(2H, m), 7.85-7.89(1H, m), 7.94-7.95(2H, m), 8.29-8.31(1H, m) (DMSO-d ₆)
1AW		-CH ₂ -CH ₂ -O-	4	H		H	O	OMe	104-105	3.12-3.20(2H, m), 3.69(3H, s), 4.39(2H, t), 4.63(2H, t), 4.71-4.75(1H, m), 6.79-6.96(5H, m), 7.15-7.26(4H, m), 7.67-7.83(3H, m), 8.17(1H, s), 8.42-8.45(1H, m)

Table 14









No.	W	A	PS	R ¹	R ²	R ³	X	Y ¹	m. p. (°C)	NMR (CDCl ₃ , δ)
1AX		-CH ₂ -CH ₂ -O-	4	H		H	O	OMe	138-140	2.58 (3H, s), 3.10-3.20 (2H, m), 3.69 (3H, s), 4.38 (2H, t), 4.59 (2H, t), 4.72-4.75 (1H, m), 6.79-6.96 (5H, m), 7.15-7.26 (4H, m), 7.73-7.83 (3H, m), 8.45-8.48 (1H, m)
1AY		-CH ₂ -CH ₂ -O-	4	H		H	O	OEt	oil	1.16 (3H, t), 3.15 (2H, d), 4.11-4.71 (7H, m), 6.76-7.23 (9H, m), 7.67-7.76 (3H, m), 8.15 (1H, s), 8.30-8.50 (1H, m)
1AZ		-CH ₂ -CH ₂ -O-	4	H		H	O	OEt	oil	1.17 (3H, t), 2.56 (3H, s), 3.15 (2H, d), 4.03-4.78 (7H, m), 6.76-7.26 (9H, m), 7.70-7.78 (3H, m), 8.40-8.50 (1H, m)
1BA		-CH ₂ -CH ₂ -O-	4	H		H	O	OH	60-62	3.19 (2H, d), 4.37 (2H, t), 4.63 (2H, t), 4.78 (1H, t), 5.50 (1H, br), 6.82-6.97 (5H, m), 7.20-7.26 (4H, m), 7.67-7.83 (3H, m), 8.18 (1H, s), 8.42-8.44 (1H, m)

Table 15







No	W	A	PS	R ¹	R ²	R ³	X	Y ¹	m. p. (°C)	NMR (CDC l ₃ , δ)
1BD		-CH ₂ -CH ₂ -O-	4	H		H	O	OH	155-159	2.57(3H, s), 3.18(2H, d), 4.35(2H, t), 4.58(2H, t), 4.76(1H, t), 6.82-6.95(5H, m), 7.16-7.26(4H, m), 7.72-7.80(3H, m), 8.44-8.46(1H, m)
1BC		-CH ₂ -CH ₂ -O-	4	H	CH ₂ CH ₃	H	O	OH	107-109	1.15(3H, t), 2.59(3H, s), 2.89-3.06(2H, m), 3.36-3.60(2H, m), 4.01-4.03(1H, m), 4.38(2H, t), 4.60(2H, t), 6.84-6.87(2H, m), 7.11-7.14(2H, m), 7.74-7.84(3H, m), 8.46-8.48(1H, m)
1BD		-CH ₂ -CH ₂ -O-	4	H	CH ₂ CH ₃	H	O	OH	114-118	1.14(3H, t), 2.88-3.06(2H, m), 3.36-3.61(2H, m), 3.99-4.02(1H, m), 4.44(2H, t), 4.71(2H, t), 6.85(2H, d), 7.12(2H, d), 7.52-7.59(5H, m), 7.72-7.81(3H, m), 8.53-8.55(1H, m)
1BE		-CH ₂ -CH ₂ -O-	4	COOEt		H	O	OH	87-88	1.14(3H, t), 2.61(3H, s), 3.56(2H, s), 4.16(4H, q), 4.39(2H, t), 4.61(2H, t), 6.82-6.84(2H, m), 6.97-7.04(3H, m), 7.09-7.11(2H, m), 7.24-7.29(2H, m), 7.77-7.84(3H, m), 8.48-8.50(1H, m)

Table 16






No.	W	A	PS	R ¹	R ²	R ³	X	Y ¹	m.p. (°C)	NMR (CDCl ₃ , δ)
IBF		-CH ₂ -CH ₂ -O-	4	COOEt		H	O	OH	70-73	1.04 (3H, t), 2.59 (3H, s), 3.53 (2H, s), 4.09-4.18 (2H, m), 4.35 (2H, t), 4.57 (2H, t), 6.30 (1H, br), 6.73-7.26 (9H, m), 7.73-7.83 (3H, m), 8.42-8.44 (1H, m)
IBG		-CH ₂ -CH ₂ -O-	3	H	CH ₂ CH ₃	H	O	OH	120-122	1.16 (3H, t), 2.61 (3H, s), 2.95-3.09 (2H, m), 3.41-3.62 (2H, m), 4.06-4.09 (1H, m), 4.41-4.44 (2H, m), 4.57-4.61 (2H, m), 6.80-6.86 (3H, m), 7.15-7.19 (1H, m), 7.76-7.85 (3H, m), 8.46-8.48 (1H, m)
IBH		-CH ₂ -CH ₂ -O-	3	H	CH ₂ CH ₃	H	O	NH ₂	91-92	1.11 (3H, t), 2.60 (3H, s), 2.83-2.88 (1H, m), 3.07-3.11 (1H, m), 3.36-3.53 (2H, m), 3.90-3.93 (1H, m), 4.40 (2H, t), 4.60 (2H, t), 5.46 (1H, br), 6.48 (1H, br), 6.79-6.84 (3H, m), 7.14-7.18 (1H, m), 7.75-7.84 (1H, m), 8.46-8.48 (1H, m)
IBI		-CH ₂ -CH ₂ -O-	4	H	CH ₂ CH ₃	H	O	OEt	54-56	1.15 (3H, t), 1.21 (3H, t), 2.60 (3H, s), 2.91-2.93 (2H, m), 3.31-3.35 (1H, m), 3.56-3.60 (1H, m), 3.92-3.96 (1H, m), 4.15 (2H, q), 4.39 (2H, t), 4.60 (2H, t), 6.85 (2H, d), 7.12 (2H, d), 7.75-7.84 (3H, m), 8.46-8.48 (1H, m)

Table 17





No.	W	A	PS	R ¹	R ²	R ³	X	Y ¹	m.p. (°C)	NMR (CDCl ₃ , δ)
1BJ		-CH ₂ -CH ₂ -O-	4	H	CH ₂ CH ₃	H	O	OH	152-154	2.60 (3H, s), 3.00-3.11 (2H, m), 3.67-3.72 (1H, m), 3.95-4.00 (1H, m), 4.17-4.20 (1H, m), 4.39 (2H, t), 4.60 (2H, t), 6.85 (2H, d), 7.13 (2H, d), 7.76-7.83 (3H, m), 8.46-8.48 (1H, m)
1BK		-CH ₂ -CH ₂ -O-	4	H	CH ₂ CH ₂ OCH ₃	H	O	OH	oil	2.60 (3H, s), 2.84-2.90 (1H, m), 3.12-3.17 (1H, m), 3.39 (3H, s), 3.39-3.55 (4H, m), 4.02-4.05 (1H, m), 4.39 (2H, t), 4.60 (2H, t), 6.86 (2H, d), 7.14 (2H, d), 7.76-7.83 (3H, m), 8.46-8.48 (1H, m)
1BL		-CH ₂ -CH ₂ -O-	4	H	CH ₂ CH ₃	H	O	OH	amorphous	1.16 (3H, t), 2.90-3.08 (2H, m), 3.41-3.47 (1H, m), 3.55-3.59 (1H, m), 3.57 (3H, s), 3.99 (3H, s), 3.99-4.04 (1H, m), 4.21 (2H, t), 4.45 (2H, t), 6.85 (2H, d), 7.13 (2H, d), 7.55 (1H, s)
1BM		-CH ₂ -CH ₂ -O-	4	H	CH ₂ CH ₂ OCH ₃	H	O	OMe	67-68	2.59 (3H, s), 2.94-2.96 (2H, m), 3.29 (3H, s), 3.45-3.49 (3H, m), 3.66-3.68 (1H, m), 3.68 (3H, s), 4.04-4.08 (1H, m), 4.38 (2H, t), 4.60 (2H, t), 6.85 (2H, d), 7.06 (2H, d), 7.75-7.82 (3H, m), 8.46-8.48 (1H, m)

Table 18





No.	W	A	PS	R ¹	R ²	R ³	X	Y ¹	m.p. (°C)	NMR (CDCl ₃ , δ)
1BN		-CH ₂ -CH ₂ -O-	4	H	CH ₂ CF ₃	II	O	OMe	111-112	2.59(3H, s), 2.93-3.06(2H, m), 3.62-3.72(1H, m), 3.72(3H, s), 3.93-4.03(1H, m), 4.13-4.16(1H, m), 4.39(2H, t), 4.60(2H, t), 6.85(2H, d), 7.10(2H, d), 7.74-7.83(3H, m), 8.46-8.48(1H, m)
1BO		-CH ₂ -CH ₂ -O-	4	H	CH ₂ CH ₃	II	O	OMe	88-89	1.15(3H, t), 2.93(2H, d), 3.23-3.58(2H, m), 3.58(3H, s), 3.69(3H, s), 3.89-4.09(1H, m), 4.00(3H, s), 4.05-4.55(4H, m), 6.82(2H, d), 7.12(2H, d), 7.52(1H, s)
1BP		-CH ₂ -CH ₂ -O-	4	H	CH ₂ CH ₃	II	O	NH ₂	oil	1.12(3H, t), 2.81-2.87(1H, m), 3.04-3.08(1H, m), 3.38-3.53(2H, m), 3.58(3H, s), 3.85-3.88(1H, m), 3.99(3H, s), 4.21(2H, t), 4.44(2H, t), 5.58(1H, br), 6.43(1H, br), 6.84(2H, d), 7.13(2H, d), 7.52(1H, s)
1BQ		-CH ₂ -CH ₂ -O-	4	H	CH ₂ CH ₃	II	O	OBt	oil	1.06-1.28(6H, m), 2.92(2H, d), 3.15-3.75(2H, m), 3.86-4.26(3H, m), 4.40-4.80(4H, m), 6.79-7.17(4H, m), 7.49-7.80(8H, m), 8.45-8.60(1H, m)

Table 19







No.	W	A	PS	R ¹	R ²	R ³ X	Y ¹	m. p. (°C)	NMR (CDCl ₃ , δ)
1BR		-CH ₂ -CH ₂ -O-	4	H	CH ₂ Cl ₃	H	0	143-145	1.02 (3H, t), 2.71 (1H, dd), 2.85 (1H, dd), 3.21-3.30 (1H, m), 3.39-3.48 (1H, m), 3.73 (1H, dd), 4.25 (2H, t), 4.33 (2H, t), 6.62 (1H, d), 6.82-6.85 (2H, m), 7.03-7.11 (4H, m), 7.47-7.52 (2H, m), 7.63-7.72 (2H, m), 8.23 (1H, d) (DMSO-d ₆)
1BS		-CH ₂ -CH ₂ -O-	4	H	CH ₂ Cl ₃	H	0	79-80	1.02 (3H, t), 2.78 (1H, dd), 2.87 (1H, dd), 3.21-3.33 (1H, m), 3.46-3.57 (1H, m), 3.91 (1H, dd), 4.25 (2H, t), 4.34 (2H, t), 6.62 (1H, d), 6.82-6.86 (2H, m), 7.09-7.13 (2H, m), 7.48-7.52 (2H, m), 7.63-7.72 (2H, m), 8.24 (1H, dd), 12.25 (1H, br) (DMSO-d ₆)
1BT		-CH ₂ -CH ₂ -O-	4	H	CH ₂ CF ₃	H	0	105-106	2.99-3.04 (2H, m), 3.64-3.73 (4H, m), 3.94-4.02 (3H, m), 4.14-4.19 (3H, m), 5.36 (2H, s), 6.80 (2H, d), 6.97 (1H, dd), 7.09-7.14 (3H, m), 7.44 (1H, ddd), 7.95 (1H, dd)
1BU		-CH ₂ -CH ₂ -O-	4	H	CH ₂ Cl ₃	H	0	105-107	3.01 (1H, dd), 3.11 (1H, dd), 3.67-3.72 (1H, m), 3.93-4.00 (3H, m), 4.14-4.21 (3H, m), 5.36 (2H, s), 6.80 (2H, d), 6.97 (1H, dd), 7.10 (1H, ddd), 7.16 (2H, d), 7.44 (1H, ddd), 7.94 (1H, dd)

Table 20

No.	W	A	PS	R ¹	R ²	R ³	X	Y ¹	m. p. (°C)	NMR (CDCl ₃ , δ)
1BV		-CH ₂ -CH ₂ -O-	4	H	CH ₂ CH ₂ OC(=O)CH ₃	H	O	OH	94-96	2.90 (1H, dd), 3.16 (1H, dd), 3.37-3.47 (5H, m), 3.53-3.56 (1H, m), 3.66-3.67 (1H, m), 3.95 (2H, t), 4.04 (1H, dd), 4.18 (2H, t), 5.37 (2H, s), 6.81 (2H, d), 6.97 (1H, dd), 7.10 (1H, dd), 7.17 (2H, d), 7.44 (1H, ddd), 7.95 (1H, dd)
1BW		-CH ₂ -CH ₂ -O-	4	H	CH ₂ CH ₂ OC(=O)CH ₃	H	O	OMe	87-88	2.95-2.98 (2H, m), 3.30 (3H, s), 3.45-3.50 (3H, m), 3.67-3.71 (4H, m), 3.95 (2H, t), 4.07 (1H, dd), 4.18 (2H, t), 5.37 (2H, s), 6.79 (2H, d), 6.97 (1H, d), 7.09-7.15 (3H, m), 7.44 (1H, ddd), 7.95 (1H, dd)

[0090] The following compounds can be prepared in accordance with processes similar to the processes described in the above Preparation Examples and Examples.

2-Methoxy-3-[4-[2-(1-oxo-1,2-dihydrophthalazin-2-yl)ethoxy]phenyl]propanamide;
 2-Methylthio-3-[4-[2-(1-oxo-1,2-dihydrophthalazin-2-yl)ethoxy]phenyl]propanamide;
 2-Propylthio-3-[4-[3-(1-oxo-1,2-dihydrophthalazin-2-yl)propoxy]phenyl]propanamide;
 2-Propoxy-3-[4-[2-(1-oxo-1,2-dihydrophthalazin-2-yl)ethoxy]phenyl]propanamide;
 5 2-Propoxy-3-[4-[3-(1-oxo-1,2-dihydrophthalazin-2-yl)propoxy]phenyl]propanamide;
 2-Isopropoxy-3-[4-[2-(1-oxo-1,2-dihydrophthalazin-2-yl)ethoxy]phenyl]propanamide;
 2-Isopropoxy-3-[4-[3-(1-oxo-1,2-dihydrophthalazin-2-yl)propyl]phenyl]propanamide;
 2-Phenoxy-3-[4-[2-(1-oxo-1,2-dihydrophthalazin-2-yl)ethoxy]phenyl]propanamide;
 2-Phenylthio-3-[4-[2-(1-oxo-1,2-dihydrophthalazin-2-yl)ethoxy]phenyl]propanamide;
 10 2-Methylamino-3-[4-[2-(1-oxo-1,2-dihydrophthalazin-2-yl)ethoxy]phenyl]propanamide;
 2-Ethylamino-3-[4-[2-(1-oxo-1,2-dihydrophthalazin-2-yl)ethoxy]phenyl]propanamide;
 2-Methyl-2-propoxy-3-[4-[2-(1-oxo-1,2-dihydrophthalazin-2-yl)ethoxy]phenyl]propanamide;
 2-Methyl-2-propylthio-3-[4-[2-(1-oxo-1,2-dihydrophthalazin-2-yl)ethoxy]phenyl]propanamide;
 2-Methyl-2-propylamino-3-[4-[2-(1-oxo-1,2-dihydrophthalazin-2-yl)ethoxy]phenyl]propanamide;
 15 2-Methyl-2-isopropoxy-3-[4-[3-(1-oxo-1,2-dihydrophthalazin-2-yl)propyl]phenyl]propanamide;
 2-Ethoxy-3-[4-[2-(6-methyl-1-oxo-1,2-dihydrophthalazin-2-yl)ethoxy]phenyl]propanamide;
 2-butoxy-3-[4-[2-(4,6-dimethyl-1-oxo-1,2-dihydrophthalazin-2-yl)ethoxy]phenyl]propanamide;
 2-Ethoxy-3-[4-[2-(6-methoxy-4-phenyl-1-oxo-1,2-dihydrophthalazin-2-yl)propoxy]phenyl]propanamide;
 2-Ethylthio-3-[4-[2-(6-methoxy-4-phenyl-1-oxo-1,2-dihydrophthalazin-2-yl)propoxy]phenyl]propanamide;
 20 2-Ethylamino-3-[4-[2-(6-methoxy-4-phenyl-1-oxo-1,2-dihydrophthalazin-2-yl)propoxy]phenyl]propanamide;
 2-Ethoxy-3-[4-[2-(6-methoxy-4-(2-pyridyl)-1-oxo-1,2-dihydrophthalazin-2-yl)propoxy]phenyl]propanamide;
 2-Methylthio-3-[4-[2-(6-methoxy-4-(2-pyridyl)-1-oxo-1,2-dihydrophthalazin-2-yl)ethoxy]phenyl]propanamide;
 2-Methylamino-3-[4-[2-(6-methoxy-4-(2-pyridyl)-1-oxo-1,2-dihydrophthalazin-2-yl)ethoxy]phenyl]propanamide;
 2-Ethoxy-3-[4-[2-(6-methoxy-4-(4-pyridyl)-1-oxo-1,2-dihydrophthalazin-2-yl)ethoxy]phenyl]propanamide;
 25 2-Ethoxy-3-[4-[3-(4-ethyl-6-methoxy-1-oxo-1,2-dihydrophthalazin-2-yl)butoxy]phenyl]propanamide;
 N1-Ethyl-2-[4-[2-(1-oxo-1,2-dihydrophthalazin-2-yl)-1-methylethoxy]-3-methylbenzyl]-2-ethoxybutanamide;
 6-Chloro-2-[3-[4-(2-ethoxy-2-methyl-3-oxo-3-piperidinopropyl)phenyl]-2-methyl-3-oxopropyl]-1,2-dihydro-1-phthalazinone;
 7-Methyl-2-[3-[4-(2-ethoxy-2-methyl-3-oxo-3-pyrrolidinopropyl)phenyl]-2-hydroxypropyl]-1,2-dihydro-1-phthalazinone;
 30 Ethyl 2-methylthio-3-[4-[2-(1-oxo-1,2-dihydrophthalazin-2-yl)ethoxy]phenyl]propanoate;
 Methyl 2-isopropoxy-3-[4-[3-(1-oxo-1,2-dihydrophthalazin-2-yl)propyl]phenyl]propanoate;
 Propyl 2-phenoxy-3-[4-[2-(1-oxo-1,2-dihydrophthalazin-2-yl)ethoxy]phenyl]propanoate;
 Isopropyl 2-ethylamino-3-[4-[2-(1-oxo-1,2-dihydrophthalazin-2-yl)ethoxy]phenyl]propanoate;
 35 2-Isopropoxy-3-[4-[3-(1-oxo-1,2-dihydrophthalazin-2-yl)propyl]phenyl]propanoic acid;
 2-Isopropylthio-3-[4-[2-(1-oxo-1,2-dihydrophthalazin-2-yl)ethoxy]phenyl]propanoic acid;
 2-Isopropylamino-3-[4-[2-(1-oxo-1,2-dihydrophthalazin-2-yl)ethoxy]phenyl]propanoic acid;
 2-Isopropoxy-3-[4-[4-(1-oxo-1,2-dihydrophthalazin-2-yl)butyl]phenyl]propanoic acid;
 2-Phenoxy-3-[4-[2-(1-oxo-1,2-dihydrophthalazin-2-yl)propoxy]phenyl]propanoic acid;
 40 2-Propoxy-3-[4-[2-(5-methyl-4-oxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)ethoxy]phenyl]propanamide;
 2-Ethylthio-3-[4-[2-(5-methyl-4-oxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)ethoxy]phenyl]propanamide;
 2-Ethylamino-3-[4-[2-(5-methyl-4-oxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)ethoxy]phenyl]propanamide;
 2-Ethoxy-3-[4-[2-(6-acetyl-4-oxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)ethoxy]phenyl]propanamide;
 2-Ethylthio-3-[4-[2-(6-amino-4-oxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)ethoxy]phenyl]propanamide;
 45 2-Ethylthio-3-[4-[2-(6-amino-4-oxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)ethoxy]phenyl]propanamide;
 2-Ethoxy-3-[4-[2-(6-methoxy-4-oxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)propoxy]phenyl]propanamide;
 2-Ethoxy-3-[4-[3-(7-methoxy-4-oxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)propyl]phenyl]propanamide;
 2-Ethoxy-3-[4-[3-(7-methyl-4-oxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)propyl]phenyl]propanamide;
 2-Phenoxy-3-[4-[3-(7-methyl-4-oxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)propyl]phenyl]propanamide;
 50 Ethyl 2-ethoxy-3-[4-[3-(4-oxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)propyl]phenyl]propanoate;
 Propyl 2-methoxy-3-[4-[2-(4-oxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)ethoxy]phenyl]propanoate;
 Ethyl 2-phenoxy-3-[4-[3-(4-oxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)propyl]phenyl]propanoate;
 2-Ethoxy-3-[4-[3-(7-methyl-4-oxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)propyl]phenyl]propanoic acid;
 2-Ethoxy-3-[4-[3-(7-chloro-4-oxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)propoxy]phenyl]propanoic acid;
 55 2-Butoxy-3-[4-[4-(7-chloro-4-oxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)butoxy]phenyl]propanoic acid;
 2-Benzyloxy-3-[4-[3-(7-cyano-4-oxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)butoxy]phenyl]propanoic acid;
 N1-Ethyl-3-[4-[4-(7-cyano-4-oxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)butoxy]phenyl]-2-ethoxy-2-methylpropanamide;

- N1,N1-Dimethyl-3-[4-[2-(7-benzyloxy-4-oxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)ethoxy]phenyl]-2-ethoxy-2-methylpropanamide;
 N1,N1-Diethyl-3-[4-[4-(7-benzyloxy-4-oxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)butoxy]phenyl]-2-ethoxy-2-methylpropanamide;
 5 6-Chloro-3-[3-[4-(2-ethoxy-2-methyl-3-oxo-3-piperidinopropyl)phenyl]-2-methyl-3-oxopropyl]-4-oxo-3,4-dihydro-2H-1,3-benzoxazine;
 2-Ethoxy-3-[4-[2-(6-methyl-4-oxo-1-phenyl-1,2,3,4-tetrahydroquinazolin-3-yl)ethoxy]phenyl]propanamide;
 2-Ethylthio-3-[4-[2-(6-methyl-4-oxo-1-phenyl-1,2,3,4-tetrahydroquinazolin-3-yl)ethoxy]phenyl]propanamide;
 2-Methoxy-3-[4-[2-(1,6-dimethyl-4-oxo-1,2,3,4-tetrahydroquinazolin-3-yl)ethoxy]phenyl]propanamide;
 10 2-Propylamino-3-[4-[3-(1,6-dimethyl-4-oxo-1,2,3,4-tetrahydroquinazolin-3-yl)propyl]phenyl]propanamide;
 2-Propylthio-3-[4-[3-(1,6-dimethyl-4-oxo-1,2,3,4-tetrahydroquinazolin-3-yl)propyl]phenyl]propanamide;
 2-Ethoxy-3-[4-[3-(1-ethyl-6-methoxy-4-oxo-1,2,3,4-tetrahydroquinazolin-3-yl)propyl]phenyl]propanamide;
 2-Isopropylthio-3-[4-[2-(1-ethyl-6-methoxy-4-oxo-1,2,3,4-tetrahydroquinazolin-3-yl)ethoxy]phenyl]propanamide;
 N1,N1-Dimethyl-2-[4-[2-(1-methyl-4-oxo-1,2,3,4-tetrahydroquinazolin-3-yl)ethoxy]benzyl]-2-ethoxybutanamide;
 15 N1,N1-Dimethyl-2-[4-[2-(1-methyl-4-oxo-1,2,3,4-tetrahydroquinazolin-3-yl)ethoxy]benzyl]-2-ethylthiobutanamide;
 Methyl 2-ethoxy-3-[4-[2-(6-methyl-4-oxo-1-phenyl-1,2,3,4-tetrahydroquinazolin-3-yl)ethoxy]phenyl]propanoate;
 Methyl 2-ethoxy-3-[4-[2-(1,6-dimethyl-4-oxo-1-phenyl-1,2,3,4-tetrahydroquinazolin-3-yl)ethoxy]phenyl]propanoate;
 Isopropyl 2-ethylthio-3-[4-[2-(1-ethyl-6,7-dimethyl-4-oxo-1,2,3,4-tetrahydroquinazolin-3-yl)ethoxy]phenyl]propanoate;
 20 Ethyl 2-ethylamino-3-[4-[2-(1-ethyl-6-methyl-4-oxo-1,2,3,4-tetrahydroquinazolin-3-yl)propoxy]phenyl]propanoate;
 2-Ethoxy-3-[4-[3-(1-ethyl-6-methyl-4-oxo-1,2,3,4-tetrahydroquinazolin-3-yl)propyl]phenyl]propanoic acid;
 2-Ethylthio-3-[4-[3-(1-ethyl-6-methyl-4-oxo-1,2,3,4-tetrahydroquinazolin-3-yl)propyl]phenyl]propanoic acid;
 2-Ethylthio-3-[4-[2-(1-ethyl-6,7-dimethyl-4-oxo-1,2,3,4-tetrahydroquinazolin-3-yl)ethoxy]phenyl]propanoic acid;
 2-Methoxy-3-[4-[2-(1-ethyl-7-methoxy-4-oxo-1,2,3,4-tetrahydroquinazolin-3-yl)ethoxy]phenyl]propanoic acid;
 25 2-Propoxy-3-[4-[2-(1-ethyl-8-methyl-4-oxo-1,2,3,4-tetrahydroquinazolin-3-yl)ethoxy]phenyl]propanoic acid;
 2-Isopropoxy-3-[4-[4-(1-ethyl-6-methyl-4-oxo-1,2,3,4-tetrahydroquinazolin-3-yl)butoxy]phenyl]propanoic acid;
 2-Ethylthio-3-[4-[2-(1-ethyl-6-methyl-4-oxo-1,2,3,4-tetrahydroquinazolin-3-yl)propoxy]phenyl]propanoic acid;
 2-Ethylamino-3-[4-[2-(1-ethyl-6-methyl-4-oxo-1,2,3,4-tetrahydroquinazolin-3-yl)propoxy]phenyl]propanoic acid;
 2-Ethoxy-3-[4-[4-(1,7-dimethyl-4-oxo-1,2,3,4-tetrahydroquinazolin-3-yl)butanoyl]phenyl]propanoic acid;
 30 2-Ethoxy-3-[4-[3-(1,7-dimethyl-4-oxo-1,2,3,4-tetrahydroquinazolin-3-yl)propanoyl]phenyl]propanoic acid;
 2-Ethylthio-3-[4-[4-(1,7-dimethyl-4-oxo-1,2,3,4-tetrahydroquinazolin-3-yl)butanoyl]phenyl]propanoic acid;
 2-Ethylthio-3-[4-[4-(1,7-dimethyl-4-oxo-1,2,3,4-tetrahydroquinazolin-3-yl)butanoyl]phenyl]propanoic acid;
 3-[4-[4-(2-Carboxy-2-ethoxybutyl)phenoxy]propyl]-7-methoxy-1-methyl-4-oxo-1,2,3,4-tetrahydro-6-quinazoline-carboxylic acid;
 35 2-Isopropoxy-3-[4-[2-(4-oxo-3,4-dihydro-2H-1,3-benzthiazin-3-yl)ethoxy]phenyl]propanamide;
 2-Ethylthio-3-[4-[2-(4-oxo-3,4-dihydro-2H-1,3-benzthiazin-3-yl)ethoxy]phenyl]propanamide;
 2-Ethoxy-3-[4-[2-(6-methyl-4-oxo-3,4-dihydro-2H-1,3-benzthiazin-3-yl)ethoxy]phenyl]propanamide;
 N1,2-Dimethyl-2-ethoxy-3-[4-[2-(4-oxo-3,4-dihydro-2H-1,3-benzthiazin-3-yl)ethoxy]phenyl]propanamide;
 2-Ethoxy-3-[4-[3-(4-oxo-3,4-dihydro-2H-1,3-benzthiazin-3-yl)propyl]phenyl]propanamide;
 40 2-Ethylamino-3-[4-[3-(4-oxo-3,4-dihydro-2H-1,3-benzthiazin-3-yl)propyl]phenyl]propanamide;
 2-Ethoxy-3-[4-[2-(4-oxo-3,4-dihydro-2H-1,3-benzthiazin-3-yl)propyl]phenyl]propanoic acid;
 2-Ethylthio-3-[4-[2-(4-oxo-3,4-dihydro-2H-1,3-benzthiazin-3-yl)ethoxy]phenyl]propanoic acid;
 2-Ethylamino-3-[4-[2-(4-oxo-3,4-dihydro-2H-1,3-benzthiazin-3-yl)ethoxy]phenyl]propanoic acid;
 2-Ethoxy-3-[4-[2-(1-oxo-1,2,3,4-tetrahydroisoquinolin-2-yl)ethoxy]phenyl]propanamide;
 45 2-Propoxy-3-[4-[2-(1-oxo-1,2,3,4-tetrahydroisoquinolin-2-yl)ethoxy]phenyl]propanamide;
 2-Ethoxy-3-[4-[3-(1-oxo-1,2,3,4-tetrahydroisoquinolin-2-yl)propyl]phenyl]propanamide;
 2-Ethylthio-3-[4-[2-(1-oxo-1,2,3,4-tetrahydroisoquinolin-2-yl)ethoxy]phenyl]propanamide;
 2-Ethylamino-3-[4-[2-(1-oxo-1,2,3,4-tetrahydroisoquinolin-2-yl)ethoxy]phenyl]propanamide;
 N1,2-dimethyl-2-ethoxy-3-[4-[2-(1-oxo-1,2,3,4-tetrahydroisoquinolin-2-yl)ethoxy]phenyl]propanamide;
 50 Methyl 2-ethoxy-3-[4-[2-(1-oxo-1,2,3,4-tetrahydroisoquinolin-2-yl)ethoxy]phenyl]propanoate;
 Isopropyl 2-ethoxy-3-[4-[3-(1-oxo-1,2,3,4-tetrahydroisoquinolin-2-yl)propyl]phenyl]propanoate;
 2-Ethoxy-3-[4-[2-(1-oxo-1,2,3,4-tetrahydroisoquinolin-2-yl)ethoxy]phenyl]propanoic acid;
 2-Methoxy-3-[4-[3-(1-oxo-1,2,3,4-tetrahydroisoquinolin-2-yl)propyl]phenyl]propanoic acid;
 2-Isopropoxy-3-[4-[2-(3-methyl-4-oxo-3,4-dihydroquinazolin-2-yl)ethoxy]phenyl]propanamide;
 55 2-Phenoxy-3-[4-[2-(3-methyl-4-oxo-3,4-dihydroquinazolin-2-yl)ethoxy]phenyl]propanamide;
 2-Propoxy-3-[4-[3-(3-ethyl-4-oxo-3,4-dihydroquinazolin-2-yl)propyl]phenyl]propanamide;
 2-Isopropylthio-3-[4-[2-(3-methyl-4-oxo-3,4-dihydroquinazolin-2-yl)ethoxy]phenyl]propanamide;
 2-Ethylamino-3-[4-[2-(3-methyl-4-oxo-3,4-dihydroquinazolin-2-yl)ethoxy]phenyl]propanamide;

N1,2-Dimethyl-2-ethoxy-3-{4-[2-(3-methyl-4-oxo-3,4-dihydroquinazolin-2-yl)ethoxy]phenyl}propanamide;

Ethyl 2-isopropylthio-3-{4-[2-(3-methyl-4-oxo-3,4-dihydroquinazolin-2-yl)ethoxy]phenyl}propanoate;

Isopropyl 2-ethylamino-3-{4-[2-(3-methyl-4-oxo-3,4-dihydroquinazolin-2-yl)ethoxy]phenyl}propanoate;

2-Ethoxy-3-{4-[2-(3-methyl-4-oxo-3,4-dihydroquinazolin-2-yl)ethoxy]phenyl}propanoic acid;

2-Ethoxy-3-{4-[3-(3-methyl-4-oxo-3,4-dihydroquinazolin-2-yl)propoxy]phenyl}propanoic acid;

2-Isopropylthio-3-{4-[2-(3-methyl-4-oxo-3,4-dihydroquinazolin-2-yl)ethoxy]phenyl}propanoic acid;

2-Ethylamino-3-{4-[2-(3-methyl-4-oxo-3,4-dihydroquinazolin-2-yl)ethoxy]phenyl}propanoic acid;

2-Isopropoxy-3-{4-[2-(1-ethyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-3-yl)ethoxy]phenyl}propanamide;

2-Methoxy-3-{4-[3-(1-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-3-yl)propyl]phenyl}propanamide;

2-Isopropoxy-3-{4-[2-(6,7-dimethoxy-1-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-3-yl)ethoxy]phenyl}propanamide;

2-Isopropylthio-3-{4-[2-(1-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-3-yl)ethoxy]phenyl}propanamide;

2-Isopropylamino-3-{4-[2-(1-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-3-yl)ethoxy]phenyl}propanamide;

N1,2-Dimethyl-2-ethoxy-3-{4-[2-(1-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-3-yl)ethoxy]phenyl}propanamide;

2-Isopropoxy-2-methyl-3-{4-[2-(1-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-3-yl)ethoxy]phenyl}propanamide;

Ethyl 2-isopropoxy-3-{4-[2-(1-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-3-yl)ethoxy]phenyl}propanoate;

Propyl 2-methoxy-3-{4-[3-(1-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-3-yl)propyl]phenyl}propanoate;

2-Isopropoxy-3-{4-[2-(1-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-3-yl)ethoxy]phenyl}propanoic acid;

2-Methoxy-3-{4-[3-(1-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-3-yl)propyl]phenyl}propanoic acid;

2-Isopropoxy-3-{4-[2-(6,7-dimethoxy-1-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-3-yl)ethoxy]phenyl}propanoic acid;

2-Isopropoxy-2-methyl-3-{4-[2-(1-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-3-yl)ethoxy]phenyl}propanoic acid;

2-Isopropoxy-3-{4-[2-(2,4-dioxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)ethoxy]phenyl}propanamide;

2-Propoxy-3-{4-[3-(2,4-dioxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)propyl]phenyl}propanamide;

2-Ethoxy-2-methyl-3-{4-[3-(2,4-dioxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)propyl]phenyl}propanamide;

2-Ethylthio-3-{4-[3-(2,4-dioxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)propyl]phenyl}propanamide;

2-Ethylamino-3-{4-[3-(2,4-dioxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)propyl]phenyl}propanamide;

Methyl 2-isopropoxy-3-{4-[2-(2,4-dioxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)ethoxy]phenyl}propanoate;

Ethyl 2-ethoxy-2-methyl-3-{4-[3-(2,4-dioxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)propyl]phenyl}propanoate;

2-Isopropoxy-3-{4-[2-(2,4-dioxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)ethoxy]phenyl}propanoic acid;

2-Propoxy-3-{4-[3-(2,4-dioxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)propyl]phenyl}propanoic acid;

2-Ethoxy-2-methyl-3-{4-[3-(2,4-dioxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)propyl]phenyl}propanoic acid;

2-Ethylthio-3-{4-[3-(2,4-dioxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)propyl]phenyl}propanoic acid;

2-Ethylamino-3-{4-[3-(2,4-dioxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)propyl]phenyl}propanoic acid;

2-Ethoxy-3-{4-[2-(2,4-dioxo-3,4-dihydro-2H-1,3-benzthiazin-3-yl)ethoxy]phenyl}propanamide;

2-Ethoxy-3-{4-[3-(2,4-dioxo-3,4-dihydro-2H-1,3-benzthiazin-3-yl)propyl]phenyl}propanamide;

2-Ethylthio-3-{4-[2-(2,4-dioxo-3,4-dihydro-2H-1,3-benzthiazin-3-yl)ethoxy]phenyl}propanamide;

2-Ethylamino-3-{4-[2-(2,4-dioxo-3,4-dihydro-2H-1,3-benzthiazin-3-yl)ethoxy]phenyl}propanamide;

N1,2-Dimethyl-2-ethoxy-3-{4-[2-(2,4-dioxo-3,4-dihydro-2H-1,3-benzthiazin-3-yl)ethoxy]phenyl}propanamide;

Methyl 2-ethoxy-3-{4-[2-(2,4-dioxo-3,4-dihydro-2H-1,3-benzthiazin-3-yl)ethoxy]phenyl}propanoate;

Propyl 2-ethylthio-3-{4-[2-(2,4-dioxo-3,4-dihydro-2H-1,3-benzthiazin-3-yl)ethoxy]phenyl}propanoate;

2-Ethoxy-3-{4-[2-(2,4-dioxo-3,4-dihydro-2H-1,3-benzthiazin-3-yl)ethoxy]phenyl}propanoic acid;

2-Ethoxy-3-{4-[3-(2,4-dioxo-3,4-dihydro-2H-1,3-benzthiazin-3-yl)propyl]phenyl}propanoic acid;

2-Ethylthio-3-{4-[2-(2,4-dioxo-3,4-dihydro-2H-1,3-benzthiazin-3-yl)ethoxy]phenyl}propanoic acid;

2-Ethylamino-3-{4-[2-(2,4-dioxo-3,4-dihydro-2H-1,3-benzthiazin-3-yl)ethoxy]phenyl}propanoic acid;

2-Isopropoxy-3-{4-[2-(4-oxo-3,4-dihydroquinazolin-3-yl)ethoxy]phenyl}propanamide;

2-Phenoxy-3-{4-[2-(4-oxo-3,4-dihydroquinazolin-3-yl)ethoxy]phenyl}propanamide;

2-Methyl-2-propoxy-3-{4-[3-(4-oxo-3,4-dihydroquinazolin-3-yl)propyl]phenyl}propanamide;

2-Isopropylthio-3-{4-[2-(4-oxo-3,4-dihydroquinazolin-3-yl)ethoxy]phenyl}propanamide;

2-Ethylamino-3-{4-[2-(4-oxo-3,4-dihydroquinazolin-3-yl)ethoxy]phenyl}propanamide;

N1,2-Dimethyl-3-{4-[2-(4-oxo-3,4-dihydroquinazolin-3-yl)ethoxy]phenyl}-2-ethoxypropanamide;



Methyl 2-ethoxy-3-{4-[2-(4-oxo-3,4-dihydroquinazolin-3-yl)ethoxy]phenyl}propanoate;

Ethyl 2-isopropylthio-3-{4-[2-(4-oxo-3,4-dihydroquinazolin-3-yl)ethoxy]phenyl}propanoate;

Ethyl 2-ethylamino-3-{4-[2-(4-oxo-3,4-dihydroquinazolin-3-yl)ethoxy]phenyl}propanoate;

2-Ethoxy-3-{4-[2-(4-oxo-3,4-dihydroquinazolin-3-yl)ethoxy]phenyl}propanoic acid;

2-Ethoxy-3-{4-[3-(4-oxo-3,4-dihydroquinazolin-3-yl)propyl]phenyl}propanoic acid;



- 2-Isopropylthio-3-{4-[2-(4-oxo-3,4-dihydroquinazolin-3-yl)ethoxy]phenyl}propanoic acid;
 2-Ethylamino-3-{4-[2-(4-oxo-3,4-dihydroquinazolin-3-yl)ethoxy]phenyl}propanoic acid;
 2-Methyl-2-propoxy-3-{4-[3-(4-oxo-3,4-dihydroquinazolin-3-yl)propyl]phenyl}propanoic acid;
 2-Isopropoxy-3-{4-[2-(1-oxo-1,2-dihydroisoquinolin-2-yl)ethoxy]phenyl}propanamide;
 2-Phenoxy-3-{4-[2-(1-oxo-1,2-dihydroisoquinolin-2-yl)ethoxy]phenyl}propanamide;
 2-Methyl-2-propoxy-3-{4-[3-(1-oxo-1,2-dihydroisoquinolin-2-yl)propyl]phenyl}propanamide;
 2-Isopropylthio-3-{4-[2-(1-oxo-1,2-dihydroisoquinolin-2-yl)ethoxy]phenyl}propanamide;
 2-Ethylamino-3-{4-[2-(1-oxo-1,2-dihydroisoquinolin-2-yl)ethoxy]phenyl}propanamide;
 N1,2-Dimethyl-3-{4-[2-(1-oxo-1,2-dihydroisoquinolin-2-yl)ethoxy]phenyl}-2-ethoxypropanamide;
 Methyl 2-ethoxy-3-{4-[2-(1-oxo-1,2-dihydroisoquinolin-2-yl)ethoxy]phenyl}propanoate;
 Ethyl 2-isopropylthio-3-{4-[2-(1-oxo-1,2-dihydroisoquinolin-2-yl)ethoxy]phenyl}propanoate;
 2-Ethoxy-3-{4-[3-(1-oxo-1,2-dihydroisoquinolin-2-yl)propyl]phenyl}propanoic acid;
 2-Isopropylthio-3-{4-[2-(1-oxo-1,2-dihydroisoquinolin-2-yl)ethoxy]phenyl}propanoic acid;
 2-Ethylamino-3-{4-[2-(1-oxo-1,2-dihydroisoquinolin-2-yl)ethoxy]phenyl}propanoic acid;
 2-Methyl-2-propoxy-3-{4-[3-(1-oxo-1,2-dihydroisoquinolin-2-yl)propyl]phenyl}propanoic acid;
 2-Ethoxy-3-{4-[2-(4-oxo-3,4-dihydro-1,2,3-benzotriazin-3-yl)ethoxy]phenyl}propanamide;
 2-Ethoxy-3-{4-[3-(4-oxo-3,4-dihydro-1,2,3-benzotriazin-3-yl)propyl]phenyl}propanamide;
 2-Isopropoxy-3-{4-[3-(4-oxo-3,4-dihydro-1,2,3-benzotriazin-3-yl)propyl]phenyl}propanamide;
 2-Ethylthio-3-{4-[2-(4-oxo-3,4-dihydro-1,2,3-benzotriazin-3-yl)ethoxy]phenyl}propanamide;
 2-Ethylamino-3-{4-[2-(4-oxo-3,4-dihydro-1,2,3-benzotriazin-3-yl)ethoxy]phenyl}propanamide;
 2-Ethoxy-2-methyl-3-{4-[2-(4-oxo-3,4-dihydro-1,2,3-benzotriazin-3-yl)ethoxy]phenyl}propanamide;
 N1,2-Dimethyl-3-{4-[2-(4-oxo-3,4-dihydro-1,2,3-benzotriazin-3-yl)ethoxy]phenyl}-2-ethoxypropanamide;
 Methyl 2-ethoxy-3-{4-[2-(4-oxo-3,4-dihydro-1,2,3-benzotriazin-3-yl)ethoxy]phenyl}propanoate;
 Ethyl 2-ethylthio-3-{4-[2-(4-oxo-3,4-dihydro-1,2,3-benzotriazin-3-yl)ethoxy]phenyl}propanoate;
 Ethyl 2-ethoxy-2-methyl-3-{4-[2-(4-oxo-3,4-dihydro-1,2,3-benzotriazin-3-yl)ethoxy]phenyl}propanoate;
 2-Ethoxy-3-{4-[2-(4-oxo-3,4-dihydro-1,2,3-benzotriazin-3-yl)ethoxy]phenyl}propanoic acid;
 2-Ethoxy-3-{4-[3-(4-oxo-3,4-dihydro-1,2,3-benzotriazin-3-yl)propyl]phenyl}propanoic acid;
 2-Isopropoxy-3-{4-[3-(4-oxo-3,4-dihydro-1,2,3-benzotriazin-3-yl)propyl]phenyl}propanoic acid;
 2-Ethylthio-3-{4-[2-(4-oxo-3,4-dihydro-1,2,3-benzotriazin-3-yl)ethoxy]phenyl}propanoic acid;
 2-Ethylamino-3-{4-[2-(4-oxo-3,4-dihydro-1,2,3-benzotriazin-3-yl)ethoxy]phenyl}propanoic acid;
 2-Ethoxy-2-methyl-3-{4-[2-(4-oxo-3,4-dihydro-1,2,3-benzotriazin-3-yl)ethoxy]phenyl}propanoic acid;
 2-Methoxy-3-{4-[2-(3-ethyl-4-methyl-2-oxo-2,5-dihydro-1H-1-pyrrolyl)ethoxy]phenyl}propanamide;
 2-Isopropoxy-3-{4-[2-(3,4-dimethyl-2-oxo-2,5-dihydro-1H-1-pyrrolyl)ethoxy]phenyl}propanamide;
 2-Isopropoxy-3-{4-[3-(3-ethyl-4-methyl-2-oxo-2,5-dihydro-1H-1-pyrrolyl)propyl]phenyl}propanamide;
 2-Isopropylthio-3-{4-[3-(3-ethyl-4-methyl-2-oxo-2,5-dihydro-1H-1-pyrrolyl)propyl]phenyl}propanamide;
 2-Isopropylamino-3-{4-[2-(3-ethyl-4-methyl-2-oxo-2,5-dihydro-1H-1-pyrrolyl)ethoxy]phenyl}propanamide;
 Ethyl 2-methoxy-3-{4-[2-(3-ethyl-4-methyl-2-oxo-2,5-dihydro-1H-1-pyrrolyl)ethoxy]phenyl}propanoate;
 Methyl 2-isopropylthio-3-{4-[3-(3-ethyl-4-methyl-2-oxo-2,5-dihydro-1H-1-pyrrolyl)propyl]phenyl}propanoate;
 2-Methoxy-3-{4-[2-(3-ethyl-4-methyl-2-oxo-2,5-dihydro-1H-1-pyrrolyl)ethoxy]phenyl}propanoic acid;
 2-Isopropoxy-3-{4-[2-(3,4-dimethyl-2-oxo-2,5-dihydro-1H-1-pyrrolyl)ethoxy]phenyl}propanoic acid;
 2-Isopropoxy-3-{4-[3-(3-ethyl-4-methyl-2-oxo-2,5-dihydro-1H-1-pyrrolyl)propyl]phenyl}propanoic acid;
 2-Isopropylthio-3-{4-[3-(3-ethyl-4-methyl-2-oxo-2,5-dihydro-1H-1-pyrrolyl)propyl]phenyl}propanoic acid;
 2-Isopropylamino-3-{4-[2-(3-ethyl-4-methyl-2-oxo-2,5-dihydro-1H-1-pyrrolyl)ethoxy]phenyl}propanoic acid;
 2-Ethoxy-3-{4-[2-(4-methyl-7-oxo-1,4-diazepan-1-yl)ethoxy]phenyl}propanamide;
 2-Ethoxy-3-{4-[3-(4-methyl-7-oxo-1,4-diazepan-1-yl)propyl]phenyl}propanamide;
 2-Ethoxy-3-{4-[3-(4-benzyl-7-oxo-1,4-diazepan-1-yl)propyl]phenyl}propanamide;
 2-Isopropoxy-3-{4-[3-(4-ethyl-7-oxo-1,4-diazepan-1-yl)propyl]phenyl}propanamide;
 2-Ethylthio-3-{4-[2-(4-methyl-7-oxo-1,4-diazepan-1-yl)ethoxy]phenyl}propanamide;
 2-Ethylamino-3-{4-[2-(4-methyl-7-oxo-1,4-diazepan-1-yl)ethoxy]phenyl}propanamide;
 N1,2-Dimethyl-3-{4-[2-(4-methyl-7-oxo-1,4-diazepan-1-yl)ethoxy]phenyl}-2-ethoxypropanamide;
 2-Ethoxy-3-{4-[2-(7-oxo-4-phenyl-1,4-diazepan-1-yl)ethoxy]phenyl}propanamide;
 Ethyl 2-ethoxy-3-{4-[2-(4-methyl-7-oxo-1,4-diazepan-1-yl)ethoxy]phenyl}propanoate;
 Methyl 2-ethylthio-3-{4-[2-(4-methyl-7-oxo-1,4-diazepan-1-yl)ethoxy]phenyl}propanoate;
 2-Ethoxy-3-{4-[2-(4-methyl-7-oxo-1,4-diazepan-1-yl)ethoxy]phenyl}propanoic acid;
 2-Ethoxy-3-{4-[3-(4-methyl-7-oxo-1,4-diazepan-1-yl)propyl]phenyl}propanoic acid;
 2-Ethoxy-3-{4-[3-(4-benzyl-7-oxo-1,4-diazepan-1-yl)propyl]phenyl}propanoic acid;
 2-Isopropoxy-3-{4-[3-(4-ethyl-7-oxo-1,4-diazepan-1-yl)propyl]phenyl}propanoic acid;
 2-Ethylthio-3-{4-[2-(4-methyl-7-oxo-1,4-diazepan-1-yl)ethoxy]phenyl}propanoic acid;

2-Ethylamino-3-[4-[2-(4-methyl-7-oxo-1,4-diazepan-1-yl)ethoxy]phenyl]propanoic acid;
 2-Ethoxy-2-methyl-3-[4-[2-(4-methyl-7-oxo-1,4-diazepan-1-yl)ethoxy]phenyl]propanoic acid;
 2-Ethoxy-3-[4-[2-(7-oxo-4-phenyl-1,4-diazepan-1-yl)ethoxy]phenyl]propanoic acid;
 2-Isopropylthio-3-[4-[2-(1-oxo-1,2-dihydrophthalazin-2-yl)ethoxy]phenyl]propanohydroxamic acid;
 2-Ethylamino-3-[4-[2-(1-oxo-1,2-dihydrophthalazin-2-yl)ethoxy]phenyl]propanohydroxamic acid;
 2-Propoxy-3-[4-[2-(4-oxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)ethoxy]phenyl]propanohydroxamic acid;
 2-Propylthio-3-[4-[2-(4-oxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)ethoxy]phenyl]propanohydroxamic acid;
 2-Ethoxy-3-[4-[2-(4-oxo-1,2,3,4-tetrahydroquinazolin-3-yl)ethoxy]phenyl]propanohydroxamic acid;
 2-Propoxy-3-[4-[2-(4-oxo-3,4-dihydro-2H-1,3-benzthiazin-3-yl)ethoxy]phenyl]propanohydroxamic acid;
 2-Ethoxy-3-[4-[2-(1-oxo-1,2,3,4-tetrahydroisoquinolin-2-yl)ethoxy]phenyl]propanohydroxamic acid;
 2-Isopropoxy-3-[4-[2-(4-oxo-3,4-dihydroquinazolin-3-yl)ethoxy]phenyl]propanohydroxamic acid;
 2-Ethoxycarbonyl-2-methoxy-3-[4-[2-(1-oxo-1,2-dihydrophthalazin-2-yl)ethoxy]phenyl]propanamide;
 N-Methyl-2-ethoxy-2-ethoxycarbonyl-3-[4-[2-(1-oxo-1,2-dihydrophthalazin-2-yl)ethoxy]phenyl]propanamide;
 2-Propoxy-3-[4-[2-(4-oxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)ethoxy]benzyl]malonamide;
 Dimethyl 2-methoxy-3-[4-[2-(1-oxo-1,2-dihydrophthalazin-2-yl)ethoxy]benzyl]malonate;
 Diethyl 2-propylthio-3-[4-[2-(4-oxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)ethoxy]benzyl]malonate;
 Diethyl 2-ethoxy-3-[4-[2-(4-oxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)ethoxy]benzyl]malonate;
 2-Ethoxycarbonyl-2-methoxy-3-[4-[2-(4-oxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)ethoxy]phenyl]propanamide;
 N-Methyl-2-ethoxy-2-ethoxycarbonyl-3-[4-[2-(4-oxo-3,4-dihydro-2H-1,3-benzoxazin-3-yl)ethoxy]phenyl]propanamide;
 2-Ethoxycarbonyl-2-isopropoxy-3-[4-[2-(4-oxo-3,4-dihydroquinazolin-3-yl)ethoxy]phenyl]propanamide;
 Diethyl 2-ethoxy-3-[4-[2-(4-oxo-3,4-dihydroquinazolin-3-yl)ethoxy]benzyl]malonate;
 2-Ethoxycarbonyl-2-methoxy-3-[4-[2-(4-oxo-3,4-dihydroquinazolin-3-yl)ethoxy]phenyl]propanamide;
 2-Ethoxy-3-[4-[2-(4-oxo-3,4-dihydroquinazolin-3-yl)ethoxy]benzyl]malonamide;
 N-Methyl-2-ethoxy-2-ethoxycarbonyl-3-[4-[2-(4-oxo-3,4-dihydroquinazolin-3-yl)ethoxy]phenyl]propanamide;
 2-Ethoxycarbonyl-2-isopropoxy-3-[4-[2-(1-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-3-yl)ethoxy]phenyl]propanamide;
 Diethyl 2-ethoxy-3-[4-[2-(1-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-3-yl)ethoxy]benzyl]malonate;
 2-Ethoxy-3-[4-[2-(1-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-3-yl)ethoxy]benzyl]malonamide; and
 N-Methyl-2-ethoxy-2-ethoxycarbonyl-3-[4-[2-(1-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-3-yl)ethoxy]phenyl]propanamide.

Test Example 1:

Antidiabetic effect on diabetic model mice:

[0091] The α -substituted phenylpropionic acid derivatives (1) according to the present invention were used to conduct a test for evaluating their activities against model mice suffering from non-insulin dependent diabetes mellitus (NIDDM). Incidentally, pioglitazone was used as a comparative compound.

(Testing method)

[0092] db/db Mice (Crea Co.; preliminarily rearing male mice aged 7 to 9 weeks for a week and using 4 or 5 mice having a blood glucose level of at least 300 mg/dl as a group) were used as subject animals. A predetermined amount of each of the compounds according to the present invention or the comparative compound was orally administered to the mice for 4 days once a day with the compound suspended in a 0.5% aqueous solution of sodium carboxymethyl cellulose. At 18 to 24 hours after the final administration, blood was collected from each of the subject animals to determine its blood glucose level (found value) by means of a simplified blood glucose meter, Mediace (manufactured by Terumo Corporation). On the other hand, the blood glucose level of a group of mice (control group) to which only a 0.5% aqueous solution of sodium carboxymethyl cellulose had been administered was also determined (control value).

[0093] On the basis of the thus-determined blood glucose level as to each of the compounds according to the present invention and the comparative compound, percent reduction of blood glucose was calculated out in accordance with the following equation. The results are shown in the following Table 21.

$$\text{Percent reduction (\%)} = [(\text{Control value} - \text{Found value}) / (\text{Control value})] \times 100$$


Table 21

Compound	Dose (mg/kg/day)	Percent reduction (%)	Compound	Dose (mg/kg/day)	Percent reduction (%)
1A	50	73	1W	50	50
1B	20	43	1X	50	60
1J	20	73	1AA	20	48
1K	50	69	1AR	20	38
1N	50	72	1BC	10	50
1O	50	42	1BI	20	52
1P	50	66	1BS	20	60
1Q	50	49	Pioglitazone	20	26
1T	50	53	Pioglitazone	50	44

[0094] As apparent from the results shown in the table, the compounds according to the present invention have an excellent antidiabetic effect.


Claims

1. An α -substituted phenylpropionic acid derivative represented by the following general formula (1):




wherein W is a monocyclic or bicyclic lactam ring which may be substituted, A is an alkylene, alkyleneoxy or alkylencarbonyl group which may be substituted by at least one hydroxy group, X is O, S, NH or CH₂, Y¹ is an amino, hydroxyamino, hydroxyalkylamino, monoalkylamino, dialkylamino, cyclic amino, hydroxy or lower alkoxy group, R¹ is a hydrogen atom, lower alkyl group, hydroxyalkyl group, alkoxyalkyl group, halogenoalkyl group or COY² (in which Y² is an amino, hydroxyamino, hydroxyalkylamino, monoalkylamino, dialkylamino, cyclic amino, hydroxy or lower alkoxy group), R² is a lower alkyl, hydroxyalkyl, alkoxyalkyl or halogenoalkyl group, COY² (in which Y² has the same meaning as defined above), or a phenyl, pyridyl or aralkyl group which may be substituted, and R³ is a hydrogen or halogen atom, or an alkyl, alkoxy, halogenoalkyl, amino, hydroxy or acyl group, or a salt thereof.


2. The α -substituted phenylpropionic acid derivative or the salt thereof according to Claim 1, wherein W in the general formula (1) is selected from among groups represented by the following (W-1) to (W-9):




(W - 1)




(W - 2)




(W - 3)




(W - 4)




(W - 5)




(W - 6)



(W - 7)



(W - 8)



(W - 9)

wherein R^4 is a hydrogen or halogen atom, an alkyl, alkoxy, halogenoalkyl, amino, hydroxy, cyano, carbamoyl, acyl, nitro, carboxy or sulfonamide group, or a phenyl or benzyloxy which may be substituted, R^5 is a hydrogen atom, an alkyl group, or an aryl, aralkyl or pyridyl group which may be substituted, R^6 is a hydrogen atom or a lower alkyl group, R^7 is a lower alkyl, phenyl or aralkyl group, Z^1 is O, S, CH_2 or NR^5 (in which R^5 has the same meaning as defined above), Z^2 is N or CH, and m is an integer of 1 to 4.

3. A medicine comprising the α -substituted phenylpropionic acid derivative or the salt thereof according to Claim 1 or 2 as an active ingredient.
4. The medicine according to Claim 3, which is an antidiabetic.
5. The medicine according to Claim 3, which is an agent for lowering lipid.
6. A medicinal composition comprising the α -substituted phenylpropionic acid derivative or the salt thereof according to Claim 1 or 2 and a pharmaceutically acceptable carrier.
7. Use of the α -substituted phenylpropionic acid derivative or the salt thereof according to Claim 1 or 2 for a medicine.



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 98 11 7153

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
A	EP 0 779 279 A (BAYER) 18 June 1997 * the whole document *	1-7	C07D237/32 C07D239/88 C07D265/22
A	WO 97 31907 A (GLAXO) 4 September 1997 * page 125; claims; example 124 *	1-7	C07D401/04 C07D239/96 C07D243/08
D,A	WO 91 19702 A (PFIZER) 26 December 1991 * the whole document *	1-7	C07D473/10 A61K31/495
D,A	CHEMICAL ABSTRACTS, vol. 126, no. 9, 3 March 1997 Columbus, Ohio, US; abstract no. 117964d, page 589; XP002032261 * abstract * & JP 08 325264 A (SUMITOMO)	1-7	
			TECHNICAL FIELDS SEARCHED (Int.Cl.6)
			C07D A61K
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 18 December 1998	Examiner Francois, J
CATEGORY OF CITED DOCUMENTS		T: theory or principle underlying the invention E: earlier patent document, but published on, or after the filing date D: document cited in the application L: document cited for other reasons &: member of the same patent family, corresponding document	
X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document			

EPO FORM 1503 03 82 (P04C01)

ANNEX TO THE EUROPEAN SEARCH REPORT ON EUROPEAN PATENT APPLICATION NO.

EP 98 11 7153

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

18-12-1998

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 779279 A	18-06-1997	DE 19546918 A	19-06-1997
		JP 9183766 A	15-07-1997
		US 5811429 A	22-09-1998
WO 9731907 A	04-09-1997	AU 2093597 A	16-09-1997
		HR 970110 A	30-04-1998
WO 9119702 A	26-12-1991	US 5089514 A	18-02-1992
		AT 149156 T	15-03-1997
		AU 646052 B	03-02-1994
		AU 7995691 A	07-01-1992
		CA 2084898 A	15-12-1991
		DE 69124798 D	03-04-1997
		DE 69124798 T	12-06-1997
		DK 533781 T	07-07-1997
		EP 0533781 A	31-03-1993
		ES 2098356 T	01-05-1997
		FI 925640 A	11-12-1992
		GR 3022714 T	30-06-1997
		HU 65603 A	28-07-1994
		IL 98447 A	31-12-1995
		JP 2581523 B	12-02-1997
		JP 7149636 A	13-06-1995
		JP 7005513 B	25-01-1995
		PT 97950 A	31-03-1992
		US 5438074 A	01-08-1995
		US 5306726 A	26-04-1994

EPO FORM P-469

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☒ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.